

2025 1 Tier Standard - SCHA
2025 Prior Authorization Criteria
CURRENT AS OF 11/01/2025

ACITRETIN

Products Affected

- *acitretin*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be a dermatologist or an oncologist.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	For prophylaxis of skin cancer in patients with previously treated skin cancers who have undergone an organ transplantation the request will be approved. For psoriasis: the patient has documented trial of, contraindication to, or medical reason for not using at least 2 of the treatment options listed: topical steroids, tazarotene, methotrexate, and cyclosporine.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

ACL INHIBITORS

Products Affected

- NEXLETOLE
- NEXLIZET

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	18 years or older
Prescriber Restrictions	Prescriber must be a cardiologist or specialist in the treatment of lipid disorders
Coverage Duration	New starts will be authorized for 4 months. Cont of therapy or reauth until end of contract year.
Other Criteria	For new starts ALL of the following must be provided: 1) Documentation of baseline low density lipoprotein cholesterol (LDL-C), 2) Member has tried and failed a high-intensity statin (i.e. atorvastatin 40-80 mg, rosuvastatin 20-40 mg) at maximum tolerated dose for 3 months via claim history or chart notes OR documentation has been provided that the member is not able to tolerate a statin. In addition to the initial criteria above if the new start is for the diagnosis of hyperlipidemia, the following are required: 1) Member has a diagnosis of heterozygous familial hypercholesterolemia (FH) OR primary hyperlipidemia, 2) Member has tried ezetimibe at a maximum tolerated dose and LDL-C is not at goal, or documentation has been provided that the patient is not able to tolerate ezetimibe. In addition to the initial criteria above if the new start is for cardiovascular risk reduction, the following are required: 1) Member has established cardiovascular disease (documented history of coronary artery disease, symptomatic peripheral artery disease, and/or cerebrovascular atherosclerotic disease, 2) Member does not have established cardiovascular disease but is considered high risk (one of the following): Diabetes Mellitus (Type 1 or Type 2) in females over 65 years of age or males over 60 years of age OR a Reynolds Risk score greater than 30% or a SCORE Risk score greater than 7.5% over 10 years OR a coronary artery calcium score greater than 400 Agaston units at any time in the past, 3) Member has a fasting LDL-C greater than or equal to 70 mg/dL. For

PA Criteria	Criteria Details
	continuation of therapy or reauthorization requests for all indications: Documentation provided that the member has obtained clinical benefit from medication (e.g., LDL-C lowering from baseline)
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

ACTEMRA

Products Affected

- ACTEMRA ACTPEN
- ACTEMRA SUBCUTANEOUS

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Specialist for submitted diagnosis.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	For pJIA: Either 1) Trial of, medical reason for not using, or contraindication to 2 of the following: Enbrel, Humira, Hadlima, or Xeljanz or 2) If utilized within the past 120 days, approve for continuation of therapy. For RA: Either 1) Trial of, medical reason for not using, or contraindication to 2 of the following: Enbrel, Humira, Hadlima, Rinvoq, or Xeljanz or 2) If utilized within the past 120 days, approve for continuation of therapy. For sJIA, Giant Cell Arteritis and Systemic Sclerosis-Associated Interstitial Lung Disease: Approve
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

ACTHAR

Products Affected

- ACTHAR
- ACTHAR GEL SUBCUTANEOUS PEN-INJECTOR

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	New starts for MS exacerbation, rheumatic disorders, collagen diseases, dermatologic diseases, serum sickness, edematous state (e.g. nephrotic syndrome without uremia), and respiratory diseases: trial of, contraindication to, or medical reason for not using 1) oral corticosteroids AND 2) Cortrophin. New starts for ophthalmic disease: trial of, contraindication to, or medical reason for not using 1) oral or ophthalmic corticosteroids AND 2) Cortrophin. Continuation of therapy or reauthorization for MS exacerbation: documentation of symptom improvement and current use of a multiple sclerosis disease modifying agent for maintenance therapy. Continuation of therapy or reauthorization for all other conditions: documented evidence of response to treatment and symptom improvement.
Age Restrictions	N/A
Prescriber Restrictions	Specialist for submitted diagnosis.
Coverage Duration	MS exacerbation: 1 month. Other conditions: new start for 3 months and reauth end of contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

ADEMPAS

Products Affected

- ADEMPAS

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with PDE inhibitor or nitrate therapy
Required Medical Information	Documentation of pulmonary arterial hypertension (PAH) WHO Group I and IV classification and PAH Functional Class. Reviewer will verify available patient claim history to confirm patient is not using PDE inhibitors or nitrates.
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be a pulmonologist or cardiologist.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

ALPHA-1 PROTEINASE INHIBITORS

Products Affected

- ARALAST NP INTRAVENOUS SOLUTION RECONSTITUTED 1000 MG, 500 MG
- GLASSIA
- PROLASTIN-C INTRAVENOUS SOLUTION
- ZEMAIRA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Documentation of hereditary alpha1-antitrypsin deficiency as evident by pretreatment serum AAT levels below 11 micromol/L and progressive FEV1 or FVC decline demonstrating symptomatic lung disease.
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be a pulmonologist.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	If the medication request is for Glassia or Aralast NP, the patient has a documented medical reason (such as trial, intolerance or contraindication) for not using Prolastin-C or Zemaira to treat their medical condition.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

ALYFTREK

Products Affected

- ALYFTREK

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent use with Trikafta, Kalydeco, Orkambi, or Symdeko. Patients with unknown CFTR gene mutations.
Required Medical Information	Documentation of CFTR gene that is responsive to vanzacaftor-tezacaftor-deutivacaftor treatment.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a pulmonologist or a provider who specializes in treatment of CF.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	Cystic Fibrosis (CF) - Initial: patient must have at least one F508del mutation in the CFTR gene or a mutation in the CFTR gene that is responsive to the requested medication. Continuation of therapy: approve.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

AMBRISENTAN

Products Affected

- *ambrisentan*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Documentation of pulmonary arterial hypertension (PAH) WHO Group I classification and PAH Functional Class.
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be a pulmonologist or cardiologist.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

APOMORPHINE

Products Affected

- *apomorphine hcl subcutaneous*

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with serotonin 5-HT3 receptor antagonists.
Required Medical Information	Reviewer will verify available patient claim history to confirm patient is not using 5-HT3 receptor antagonists.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a neurologist.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	If diagnosis is Parkinson's, the patient must have a documented trial of, contraindication to, or medical reason for not using two alternatives such as entacapone, tolcapone, rasagiline, selegiline, carbidopa/levodopa, bromocriptine, pramipexole or ropinirole.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

AQNEURSA

Products Affected

- AQNEURSA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Specialist for submitted diagnosis.
Coverage Duration	New starts will be authorized for 6 months. Cont of therapy or reauth until end of contract year.
Other Criteria	For new starts: 1) The member has a documented diagnosis of Niemann-Pick disease type C (NPC) AND 2) Documentation of genetic testing identifying disease-causing alleles in NPC1 or NPC2 AND 3) Documentation of disease-related neurological symptoms (e.g., developmental delay/regression, ataxia, cataplexy, seizures, motor-function decline, tremors, dysphagia) For reauthorization: Documentation that member has had positive response to therapy (e.g., improvement in neurological status, decrease in functional Scale for Assessment and Rating of Ataxia [fSARA] score).
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

ARCALYST

Products Affected

- ARCALYST

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Specialist for submitted diagnosis.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	For deficiency of interleukin-1 receptor antagonist, documented trial of, contraindication to, or medical reason for not using Kineret. For continuation of therapy or reauthorization: Documentation has been provided that patient has clinically benefited from medication.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

ARIKAYCE

Products Affected

- ARIKAYCE

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Mycobacterium avium complex (MAC): (1) Documented diagnosis of MAC lung disease as verified by failure to achieve at least 2 negative sputum cultures following 6 consecutive months of a combination antibacterial drug regimen AND (2) Provider attestation that medication is being used as part of a combination antibacterial drug regimen.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a pulmonologist or an infectious disease specialist
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

ARISTADA

Products Affected

- ARISTADA INITIO 441 MG/1.6ML, 662 MG/2.4ML, 882
- ARISTADA INTRAMUSCULAR MG/3.2ML
- PREFILLED SYRINGE 1064 MG/3.9ML,

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	The member has a documented history of receiving oral aripiprazole without any clinically significant side effects.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	Trial of, contraindication to, or medical reason (e.g. intolerance, hypersensitivity or contraindication) for not using at least two of the following: Abilify Maintena, Abilify Asimtufii, Invega Sustenna, Invega Trinza, or Risperdion Microsphere ER.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

AUVELITY

Products Affected

- AUVELITY

PA Criteria	Criteria Details
Exclusion Criteria	Seizure disorder
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	Trial of, contraindication to, or medical reason for not using to two generic antidepressants.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

AZTREONAM LYSINE

Products Affected

- CAYSTON

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be a pulmonologist, infectious disease specialist, or an expert in the treatment of cystic fibrosis.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

BENLYSTA

Products Affected

- BENLYSTA SUBCUTANEOUS

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be a rheumatologist, nephrologist, or specialist in the treatment of autoimmune disorders.
Coverage Duration	New starts will be authorized for 6 months. Cont of therapy or reauth until end of contract year.
Other Criteria	For new starts for systemic lupus erythematosus (SLE): concurrent use of two of the following or medical reason for not using glucocorticoids, azathioprine, methotrexate, mycophenolate, or hydroxychloroquine, chloroquine, and cyclophosphamide. For continuation of therapy or reauthorization for SLE: documentation of clinical response to therapy (i.e. fewer flares that required steroid treatment, lower average daily oral prednisone dose, improved daily function either as measured through a validated functional scale or through improved daily performance documented at clinic visits, etc.) For new starts for lupus nephritis (LN): concurrent use of or medical reason for not using background immunosuppressive therapy regimen. For continuation of therapy or reauthorization for LN: Documentation of improvement in renal function (i.e. reduction in UPCR).
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

BESREMI

Products Affected

- BESREMI

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be a hematologist, oncologist, or specialist for submitted diagnosis.
Coverage Duration	The request will be authorized until the end of the contract year.
Other Criteria	Trial of, contraindication to, or medical reason for not using Pegasys
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

BORUZU

Products Affected

- BORUZU

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be an oncologist or specialist for submitted diagnosis.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

BOSENTAN

Products Affected

- *bosentan oral tablet*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Documentation of pulmonary arterial hypertension (PAH) WHO Group I classification and PAH Functional Class.
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be a pulmonologist or cardiologist.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

CAMZYOS

Products Affected

- CAMZYOS

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a cardiologist.
Coverage Duration	New starts will be authorized for 6 months. Cont of therapy or reauth until end of contract year.
Other Criteria	For all new starts, ALL of the following must be provided: 1) Diagnosis of symptomatic New York Heart Association (NYHA) class II or III obstructive hypertrophic cardiomyopathy (oHCM) AND 2) Patient has a left ventricular ejection fraction (LVEF) greater than or equal to 55% AND 3) Assessment of Valsalva left ventricular outflow tract (LVOT) gradient AND 4) Trial of, medical reason for not using or contraindication to BOTH of the following: Beta blockers (i.e. metoprolol, propranolol, atenolol) AND Non-dihydropyridine calcium channel blockers (i.e. verapamil, diltiazem) AND 5) Prescriber attests that patient is not using moderate to strong CYP2C19 or CYP3A4 inhibitors or inducers. For continuation of therapy or reauthorization, all of the following must be provided: 1) Documentation of clinical benefit as evidenced by an improvement from baseline in oHCM symptoms (i.e., improvement in fatigue, chest pain, shortness of breath, LVOT, peak oxygen consumption, etc.) OR improvement or no worsening of NYHA functional class AND 2) Member must also have a left ventricular ejection fraction (LVEF) greater than or equal to 50%.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

PA Criteria	Criteria Details
Part B Prerequisite	No

CARGLUMIC ACID

Products Affected

- *carglumic acid oral tablet soluble*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

CASPOFUNGIN

Products Affected

- *caspofungin acetate*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Documentation of a consultation with an infectious disease specialist.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

CERDELGA

Products Affected

- CERDELGA

PA Criteria	Criteria Details
Exclusion Criteria	Patients with undetermined CYP2D6 metabolizer status.
Required Medical Information	Patient's CYP2D6 metabolizer status, as determined by an FDA approved test. For reauthorization, documentation has been provided that patient has obtained clinical benefit from medication (e.g. increased platelet count, improvement in anemia, PFTs, improvement in radiographic scans, improved quality of life).
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a specialist in treatment of Gaucher's disease.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

CGRP ANTAGONISTS

Products Affected

- AIMOVIG
- EMGALITY
- EMGALITY (300 MG DOSE)
- NURTEC
- QULIPTA
- UBRELVY
- ZAVZPRET

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	New starts will be authorized for 6 months. Cont of therapy or reauth until end of contract year.
Other Criteria	For acute migraine new starts - for Ubrelvy and Nurtec requests, must have trial of, contraindication to or medical reason for not using a triptan. For Zavzpret requests, must have trial of, contraindication to or medical reason for not using Ubrelvy or Nurtec. For migraine prophylaxis new starts - 1) at least 4 migraine days per month or one or more severe migraine attacks lasting for greater than 12 hours despite use of abortive therapy (e.g. triptans or NSAIDs) and 2) trial of, contraindication to, or medical reason for not using at least two of the following agents: a beta adrenergic blocker, an anti-epileptic agent, a tricyclic antidepressant, or a serotonin-norepinephrine reuptake inhibitor. For Emgality requests for episodic cluster headache new starts - must have trial of, contraindication to, or a medical reason for not using verapamil for at least 4 weeks at minimum effective doses. For continuation of therapy or reauthorization - For acute migraine (Nurtec, Ubrelvy, Zavzpret), must show documentation of improvement in migraine symptoms (pain, photophobia, phonophobia). For migraine prevention (Nurtec, Emgality, Qulipta, Aimovig), must show documentation of improvement in migraine symptoms. For episodic cluster headache treatment, must show documentation of reduction in frequency of headaches.

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

CHOLBAM

Products Affected

- CHOLBAM

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	For new starts: Patient has documented diagnosis of either: 1) bile acid synthesis disorder due to a single enzyme defect or 2) peroxisomal disorders. For continuation of therapy or reauthorization: prescriber attests: 1) the patient has clinical improvement with therapy (i.e. liver function tests) AND 2) there is no evidence of biliary obstruction or cholestasis
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be a hepatologist, gastroenterologist, or metabolic specialist
Coverage Duration	New starts will be authorized for 3 months. Cont of therapy or reauth until end of contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

CIBINQO

Products Affected

- CIBINQO

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Specialist for submitted diagnosis.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	For atopic dermatitis: Trial of, contraindication to, or medical reason for not using Rinvoq
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

CIMZIA

Products Affected

- CIMZIA (2 SYRINGE)
- CIMZIA SUBCUTANEOUS KIT 2 X 200 MG
- CIMZIA-STARTER

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Specialist for submitted diagnosis.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	For ankylosing spondylitis: Either 1) Trial of, medical reason for not using, or contraindication to 2 of the following therapies: Enbrel, Humira, Hadlima, Rinvoq or Xeljanz or 2) If utilized within the past 120 days, approve for continuation of therapy. For Crohns Disease: Either 1) Trial of, medical reason for not using, or contraindication to 2 of the following therapies: Humira, Hadlima, Skyrizi or Stelara or 2) If utilized within the past 120 days, approve for continuation of therapy. For non-radiographic axial spondyloarthritis: approve. For psoriasis: Either 1) Trial of, medical reason for not using, or contraindication to 2 of the following therapies: Skyrizi, Tremfya, Stelara, Enbrel, Hadlima, or Humira 2) If utilized within the past 120 days, approve for continuation of therapy. For PsA: Either 1) Trial of, medical reason for not using, or contraindication to 2 of the following therapies: Stelara, Skyrizi, Tremfya, Xeljanz, Rinvoq, Enbrel, Hadlima, or Humira, or 2) If utilized within the past 120 days, approve for continuation of therapy. For RA: Either 1) Trial of, medical reason for not using, or contraindication to 2 of the following: Enbrel, Humira, Hadlima, Rinvoq or Xeljanz or 2) If utilized within the past 120 days, approve for continuation of therapy
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

Formulary ID 25406

Last Update: 10/17/2025

PA Criteria	Criteria Details
Part B Prerequisite	No

CORLANOR

Products Affected

- CORLANOR ORAL SOLUTION
- *ivabradine hcl*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	New starts for chronic heart failure must have all of the following: 1) LVEF of 35% or less 2) Sinus rhythm and have resting heart rate greater than or equal to 70 bpm. For pediatric patients with heart failure due to dilated cardiomyopathy: approve.
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be a cardiologist.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	Trial of, contraindication to, or medical reason for not receiving a beta blocker.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

CORTROPHIN

Products Affected

- CORTROPHIN
- CORTROPHIN GEL

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	New starts for MS exacerbation, rheumatic disorders, collagen diseases, dermatologic diseases, serum sickness, edematous state (e.g. nephrotic syndrome without uremia), and respiratory diseases: trial of, contraindication to, or medical reason for not using oral corticosteroids. New starts for ophthalmic disease: trial of, contraindication to, or medical reason for not using oral or ophthalmic corticosteroids. Continuation of therapy or reauthorization for MS exacerbation: documentation of symptom improvement and current use of a multiple sclerosis disease modifying agent for maintenance therapy. Continuation of therapy or reauthorization for all other conditions: documented evidence of response to treatment and symptom improvement.
Age Restrictions	N/A
Prescriber Restrictions	Specialist for submitted diagnosis.
Coverage Duration	MS exacerbation: 1 month. Other conditions: new start for 3 months and reauth end of contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

COSENTYX

Products Affected

- COSENTYX
- COSENTYX (300 MG DOSE)
- COSENTYX SENSOREADY (300 MG)
- COSENTYX SENSOREADY PEN
- COSENTYX UNOREADY

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Specialist for submitted diagnosis.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	For ankylosing spondylitis: Either 1) Trial of, medical reason for not using, or contraindication to 2 of the following therapies: Enbrel, Humira, Hadlima, Rinvoq or Xeljanz, or 2) If utilized within the past 120 days, approve for continuation of therapy. For non-radiographic axial spondyloarthritis: approve. For psoriasis: Either 1) Trial of, medical reason for not using, or contraindication to 2 of the following therapies: Stelara, Skyrizi, Tremfya, Enbrel, Hadlima, or Humira, or 2) If utilized within the past 120 days, approve for continuation of therapy. For PsA: Either 1) Trial of, medical reason for not using, or contraindication (e.g., safety concerns, not indicated for patients age) to 2 of the following therapies: Stelara, Skyrizi, Tremfya, Xeljanz, Rinvoq, Enbrel, Hadlima, or Humira, or 2) If utilized within the past 120 days, approve for continuation of therapy. For enthesitis-related arthritis: approve. For moderate to severe hidradenitis suppurativa (HS): Either 1) Trial of, medical reason for not using, or contraindication (e.g., safety concerns, not indicated for patients age) to 1 of the following therapies: Hadlima or Humira, or 2) If utilized within the past 120 days, approve for continuation of therapy.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

PA Criteria	Criteria Details
Part B Prerequisite	No

CYSTAGON

Products Affected

- CYSTAGON

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

CYSTARAN

Products Affected

- CYSTARAN

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Documentation of diagnosis for cystinosis with corneal cystine crystal accumulation.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an ophthalmologist or metabolic disease specialist.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

DALFAMPRIDINE ER

Products Affected

- *dalfampridine er*

PA Criteria	Criteria Details
Exclusion Criteria	History of seizure or moderate/severe renal impairment (CrCl less than or equal to 50 mL/min).
Required Medical Information	For new starts: 1) Attestation that creatinine clearance (CrCl) greater than 50 mL/min was confirmed prior to initiation of therapy, AND 2) Documentation has been provided that member is ambulatory (able to walk at least 25 feet) and has a documented walking impairment, AND 3) For appropriate indications, member is currently being treated with a disease modifying agent (e.g. immunomodulator, interferon, etc.) or has a medical reason why member is unable to use a disease modifying agent for their condition. For continuation of therapy or re-authorization requests: 1) Member must experience improvement in walking from baseline due to use of dalfampridine ER.
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be a neurologist.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

DEFERASIROX

Products Affected

- *deferasirox*
- *deferasirox granules*

PA Criteria	Criteria Details
Exclusion Criteria	Creatinine clearance less than 40 mL/min or platelet counts less than 50,000/mm ³ .
Required Medical Information	For all indications: platelet count greater than or equal to 50,000/mm ³ (within 30 days). For chronic iron overload due to transfusions: serum ferritin concentration greater than 1000 mcg/L (lab result within 30 days). For chronic iron overload in non-transfusion-dependent thalassemia syndromes: serum ferritin concentration greater than 300 mcg/L (lab result within 30 days).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	For deferasirox granules oral packets, the member must have medical reason for not using deferasirox tablets or oral soluble tablets.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

DEFERIPRONE

Products Affected

- *deferiprone*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	For new starts: 1) serum ferritin level above 1,000 mcg/L and absolute neutrophil count (ANC) greater than $1.5 \times 10^9/L$ within 30 days of request, and 2) Trial of, contraindication to, or medical reason for not using deferasirox tablets. For continuation of therapy or reauthorization, decrease in serum ferritin from baseline.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

DIACOMIT

Products Affected

- DIACOMIT

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be a neurologist.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	For members 2 years and older: Trial of, contraindication to, or medical reason for not using one generic anticonvulsant for appropriate indications. For members under 2 years old: Approve.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

DICHLORPHENAMIDE

Products Affected

- dichlorphenamide*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a geneticist, neurologist, or endocrinologist.
Coverage Duration	New starts will be authorized for 2 months. Cont of therapy or reauth until end of contract year.
Other Criteria	Continuation of therapy or reauthorization: documentation of clinical improvement with therapy.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

DIFICID

Products Affected

- DIFICID
- fidaxomicin*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized for 10 days.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

DIHYDROERGOTAMINE NASAL

Products Affected

- *dihydroergotamine mesylate nasal*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	For new starts: Member has a diagnosis of migraine headaches with or without aura. Prescriber attestation that it will be used for the acute treatment of migraine. For continuation of therapy or reauthorization: Documentation or provider attestation of positive clinical response (e.g., improvement in pain, photophobia, phonophobia).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Requests will be authorized for 12 weeks.
Other Criteria	Trial of, contraindication to, or medical reason (e.g. intolerance, hypersensitivity or contraindication) for not using a triptan (e.g., rizatriptan, sumatriptan).
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

DOPTELET

Products Affected

- DOPTELET
- DOPTELET SPRINKLE

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	For new starts for chronic liver disease and chronic immune thrombocytopenia (chronic ITP): documented baseline platelet count of less than 50,000/mcL.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with hematologist, hepatologist or surgeon.
Coverage Duration	For thrombocytopenia with CLD getting procedure: 5 days. For chronic ITP: remainder of contract year
Other Criteria	For chronic ITP: trial of, contraindication to, or medical reason for not using a corticosteroid. For thrombocytopenia with chronic liver disease (CLD): approve.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

DOXEPIN CREAM

Products Affected

- *doxepin hcl external*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized for 1 month.
Other Criteria	Trial of, contraindication to, or medical reason for not using a topical corticosteroid or topical calcineurin inhibitor.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

DUPIXENT

Products Affected

- DUPIXENT SUBCUTANEOUS SOLUTION AUTO-INJECTOR
- DUPIXENT SUBCUTANEOUS SOLUTION PREFILLED SYRINGE

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Chronic spontaneous urticaria (CSU) - Initial: patient has inadequate symptomatic relief despite trial of two weeks of a second-generation oral antihistamine (unless contraindicated). Continuation of therapy: patient has positive clinical response to treatment. Bullous pemphigoid - Initial: patient has had trial of, contraindication to, or medical reason for not using a topical corticosteroid, oral corticosteroid, or an immunosuppressive agent. Continuation of therapy: patient has positive clinical response to treatment.
Age Restrictions	N/A
Prescriber Restrictions	Initial therapy only: prescribed by or in consultation with specialist for submitted diagnosis.
Coverage Duration	AD - Initial: 4 months, PN & CSU - Initial: 6 months, All others: end of contract year.
Other Criteria	Atopic dermatitis (AD) in patients 6 months of age and older - Initial: (1) patient has diagnosis of moderate to severe AD, AND (2) has had trial of, contraindication to, or medical reason for not using either a topical corticosteroid or topical calcineurin inhibitor. Continuation of therapy: patient has positive clinical response to treatment. Asthma with eosinophilic phenotype - Initial: (1) patient has baseline blood eosinophil count greater than or equal to 150 cells per microliter, AND (2) asthma remains inadequately controlled despite current treatment with or medical reasons for not using BOTH (i) medium to high dose inhaled corticosteroid AND (ii) additional controller (i.e. leukotriene modifier, long acting beta-2-agonist, long acting muscarinic antagonist, and sustained released theophylline). Asthma, oral corticosteroid dependent - Initial: asthma remains inadequately controlled despite current treatment with or medical reasons for not using BOTH (i) high dose inhaled corticosteroid AND (ii) additional controller (i.e. leukotriene modifier, long acting beta-2-agonist, long acting muscarinic antagonist, and sustained released theophylline). Asthma with eosinophilic phenotype or oral corticosteroid dependent -

PA Criteria	Criteria Details
	Continuation of therapy: clinical improvement in asthma control (i.e. reduction in frequency and/or severity of exacerbations and symptoms OR reduction in daily maintenance oral corticosteroid dose). Chronic rhinosinusitis with nasal polyps (CRSwNP) - Initial: (1) Dupixent is used as add-on maintenance treatment, AND (2) patient has a trial of, contraindication to, or medical reason for not using nasal corticosteroids. Continuation of therapy: patient has positive clinical response to treatment. Prurigo nodularis (PN): attestation is provided confirming diagnosis. Eosinophilic esophagitis (EoE) - Initial: (1) diagnosis has been confirmed by esophageal biopsy AND (2) patient weighs at least 15 kilograms AND (3) patient experienced an inadequate treatment response, intolerance, or has a contraindication to a topical or oral corticosteroid (i.e. fluticasone propionate or budesonide). Continuation of therapy: patient has positive clinical response to treatment. Chronic Obstructive Pulmonary Disease (COPD) - Initial: (1) documented diagnosis of COPD with an eosinophilic phenotype, AND (2) Dupixent is used as an add-on maintenance treatment, AND (3) documentation that patient's COPD is inadequately controlled. Continuation of therapy: patient has positive clinical response to treatment.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

EGRIFTA

Products Affected

- EGRIFTA SV
- EGRIFTA WR

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Documentation of active antiretroviral therapy for at least 8 weeks.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

EMSAM

Products Affected

- EMSAM

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with SSRIs, SNRIs, clomipramine and imipramine, meperidine, tramadol, methadone, pentazocine, and propoxyphene, and the antitussive agent dextromethorphan or carbamazepine
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	For new starts: Trial of, contraindication to, or medical reason for not using two generic antidepressants.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

ENBREL

Products Affected

- ENBREL MINI
- ENBREL SUBCUTANEOUS SOLUTION 25 MG/0.5ML
- ENBREL SUBCUTANEOUS SOLUTION PREFILLED SYRINGE
- ENBREL SURECLICK SUBCUTANEOUS SOLUTION AUTO-INJECTOR

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Specialist for submitted diagnosis.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	For RA: Trial of, medical reason for not using, or contraindication to 1 disease modifying antirheumatic drug (DMARD) (methotrexate, leflunomide, or sulfasalazine). For pJIA: Trial of, medical reason for not using, or contraindication to 1 of the following DMARDs: methotrexate or leflunomide. For PsA or psoriasis: approve. For ankylosing spondylitis: Trial of, medical reason for not using, or contraindication to naproxen.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

ENDARI

Products Affected

- *l-glutamine oral packet*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Documentation that two or more painful sickle cell crises have occurred in the past 12 months.
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be a hematologist.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	Trial of, contraindication to, or medical reason for not using hydroxyurea for at least three months.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

ENTYVIO

Products Affected

- ENTYVIO PEN

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Specialist for submitted diagnosis.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	For ulcerative colitis: Either 1) Trial of, medical reason for not using, or contraindication to 2 of the following therapies: Stelara, Tremfya, Hadlima or Humira. 2) If utilized within the past 120 days, approve for continuation of therapy.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

EPIDIOLEX

Products Affected

- EPIDIOLEX

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be a neurologist.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	Trial of, contraindication to, or medical reason for not using one generic anticonvulsant for appropriate indications.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

EPRONTIA

Products Affected

- *topiramate oral solution*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	The request will be authorized until the end of the contract year.
Other Criteria	Documented trial of, contraindication to, or medical reason for not using topiramate sprinkle capsule or topiramate oral tablet.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

ERYTHROPOIETIN STIMULATING AGENTS

Products Affected

- ARANESP (ALBUMIN FREE) INJECTION SOLUTION 100 MCG/ML, 200 MCG/ML, 25 MCG/ML, 40 MCG/ML, 60 MCG/ML
- ARANESP (ALBUMIN FREE) INJECTION SOLUTION PREFILLED SYRINGE
- EPOGEN INJECTION SOLUTION 10000 UNIT/ML, 2000 UNIT/ML, 20000 UNIT/ML, 3000 UNIT/ML, 4000 UNIT/ML
- PROCIT
- RETACRIT INJECTION SOLUTION 10000 UNIT/ML, 10000 UNIT/ML(1ML), 2000 UNIT/ML, 20000 UNIT/ML, 3000 UNIT/ML, 4000 UNIT/ML, 40000 UNIT/ML

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	For new starts for all indications: Hgb within compendia range for treatment of the requested medical condition. For continuation of therapy or re-authorization: Hgb must not exceed 10 g/dL (anemia related to cancer), 11 g/dL (anemia of CKD), 12 g/dL (zidovudine-related anemia in members with HIV and ribavirin-induced anemia), 13 g/dL (elective, noncardiac, nonvascular surgery needing red blood cell allogeneic transfusion reduction).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized for 6 months.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

ERZOFRI

Products Affected

- ERZOFRI INTRAMUSCULAR MG/1.5ML, 351 MG/2.25ML, 39
SUSPENSION PREFILLED SYRINGE MG/0.25ML, 78 MG/0.5ML
117 MG/0.75ML, 156 MG/ML, 234

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	The member has a documented history of receiving oral paliperidone or oral risperidone without any clinically significant side effects.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	For schizophrenia: trial and failure of, contraindication, or medical reason (e.g. intolerance, hypersensitivity or contraindication) for not using at least two of the following: Abilify Maintena, Abilify Asimtufii, Risperidone Microsphere, Invega Sustenna, Invega Trinza, and Invega Hafyera. For schizoaffective disorder: trial and failure of, contraindication, or medical reason (e.g. intolerance, hypersensitivity or contraindication) for not using Invega Sustenna.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

EUCRISA

Products Affected

- EUCRISA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be a dermatologist, immunologist or an allergist.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	For patients 2 years of age and older: Trial of, contraindication to, or medical reason for not using topical tacrolimus or pimecrolimus. For patients less than 2 years of age: approve.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

EVRYSDI

Products Affected

- EVRYSDI

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	For new starts, all of the following must be included: 1) Documentation of genetic testing confirming diagnosis AND 2) Documentation of baseline motor function or motor milestone achievement [e.g. CHOP Infant Test of Neuromuscular Disorders (CHOP-INTEND) or Hammersmith Infant Neurological Examination (HINE) for Type 1 or Hammersmith Functional Motor Scale Expanded Scores (HFMSE) for Type II and Type III, or 6 minute walk test in subjects able to walk]. For continuation of therapy or reauthorization, documentation of clinical response has been submitted (e.g. improvement in motor function/motor milestone achievement scores using CHOP-INTEND or HFMSE, 6 minute walk test or HINE improvement in more categories of motor milestones than worsening).
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be a neurologist.
Coverage Duration	New starts will be authorized for 6 months. Cont of therapy or reauth until end of contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

FABHALTA

Products Affected

- FABHALTA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be a hematologist, nephrologist, or oncologist.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

FASENRA

Products Affected

- FASENRA
- FASENRA PEN

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Specialist for submitted diagnosis.
Coverage Duration	New starts will be authorized for 6 months. Cont of therapy or reauth until end of contract year.
Other Criteria	New starts for severe asthma with an eosinophilic phenotype: 1)Baseline blood eosinophil count greater than or equal to 150 cells per microliter AND 2) symptoms persist with at least 1 exacerbation in the last 12 months requiring additional treatment (e.g. oral systemic steroids) while on a high dose inhaled corticosteroid with an additional controller medication (ie. long-acting B2 agonist). Continuation of therapy or re-authorization for severe asthma with an eosinophilic phenotype: clinical benefit from use of the drug. New starts for eosinophilic granulomatosis with polyangiitis (EGPA): trial of, contraindication to, or medical reason for not using one of the following medications: cyclophosphamide or methotrexate. Continuation of therapy or re-authorization for EGPA: clinical benefit from use of the drug.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

FENTANYL CITRATE TRANSMUCOSAL PRODUCTS

Products Affected

- *fentanyl citrate buccal lozenge on a handle*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Documentation must be provided for the all of the following: 1) fentanyl citrate oral transmucosal is being prescribed to treat cancer-related breakthrough pain AND 2) Patient has been taking opioids at a dose equal to 60 MME per day for at least one week.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

FILSPARI

Products Affected

- FILSPARI

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with renin-angiotensin-aldosterone system (RAAS) inhibitors, endothelin receptor antagonists, or aliskiren
Required Medical Information	For new starts: Attestation that member has diagnosis of primary immunoglobulin A nephropathy (IgAN) at risk of rapid disease progression. Member has an estimated glomerular filtration rate (eGFR) greater than or equal to 30 mL/min/1.73 m(2) and proteinuria. For continuation of therapy or reauthorization: Documentation of positive clinical response (ie. decrease in urine protein-to-creatinine ratio (UPCR)).
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a nephrologist.
Coverage Duration	New starts will be authorized for 9 months. Cont of therapy or reauth until end of contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

FINTEPLA

Products Affected

- FINTEPLA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be a neurologist.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	Trial of, contraindication to, or medical reason for not using one generic anticonvulsant for appropriate indications.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

FIRDAPSE

Products Affected

- FIRDAPSE

PA Criteria	Criteria Details
Exclusion Criteria	History of seizures.
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be a neurologist.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

FLUCYTOSINE

Products Affected

- *flucytosine oral*

PA Criteria	Criteria Details
Exclusion Criteria	Complete dihydropyrimidine dehydrogenase (DPD) enzyme deficiency
Required Medical Information	Attestation member is taking in combination with amphotericin B.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

FLUOROURACIL

Products Affected

- *fluorouracil external cream 0.5 %*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a dermatologist or oncologist.
Coverage Duration	Request will be authorized for 12 weeks.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

GALAFOLD

Products Affected

- GALAFOLD

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescribed must be a geneticist, cardiologist, nephrologist or specialist experienced in the treatment of Fabry disease.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

GATTEX

Products Affected

- GATTEX

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	For new starts: attestation of 1) Colonoscopy of full colon with removal of polyps within six months prior to starting treatment for adults or 2) Fecal occult blood testing within six months prior to starting treatment for pediatric patients. For continuation of therapy or reauthorization: Documentation is provided that the member has obtained a clinical benefit (e.g. reduction in parenteral fluid volume, reduction in number of days receiving parenteral nutrition).
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a gastroenterologist.
Coverage Duration	New starts will be authorized for 6 months. Cont of therapy or reauth until end of contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

GLP-1 AGONISTS

Products Affected

- MOUNJARO SUBCUTANEOUS SOLUTION AUTO-INJECTOR
- OZEMPIC (0.25 OR 0.5 MG/DOSE) SUBCUTANEOUS SOLUTION PEN-INJECTOR 2 MG/3ML
- OZEMPIC (1 MG/DOSE) SUBCUTANEOUS SOLUTION PEN-INJECTOR 4 MG/3ML
- OZEMPIC (2 MG/DOSE)
- RYBELSUS
- RYBELSUS (FORMULATION R2)
- TRULICITY SUBCUTANEOUS SOLUTION AUTO-INJECTOR
- VICTOZA SUBCUTANEOUS SOLUTION PEN-INJECTOR

PA Criteria	Criteria Details
Exclusion Criteria	The member has an indication of weight loss/obesity only or type 1 diabetes.
Required Medical Information	The member has a diagnosis of type 2 diabetes.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

GNRH AGONISTS

Products Affected

- CAMCEVI
- ELIGARD
- FIRMAGON (240 MG DOSE)
- FIRMAGON SUBCUTANEOUS SOLUTION RECONSTITUTED 80 MG
- *leuprolide acetate (3 month)*
- LUPRON DEPOT (1-MONTH)
- LUPRON DEPOT (3-MONTH)
- LUPRON DEPOT (4-MONTH)
- LUPRON DEPOT (6-MONTH)
- LUTRATE DEPOT
- TRELSTAR MIXJECT

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	If the medication request is for the treatment of prostate cancer and if the request is for any other GnRH agonist other than Eligard or leuprolide, the patient must have a documented trial of, contraindication to, or medical reason for not using Eligard or leuprolide to treat their prostate cancer.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

GOCOVRI

Products Affected

- GOCOVRI

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a neurologist.
Coverage Duration	New starts will be authorized for 3 months. Cont of therapy or reauth until end of contract year.
Other Criteria	New starts: trial of, contraindication to, or medical reason for not using generic amantadine. Continuation of therapy or reauthorization: Member demonstrates clinical benefit (i.e. improvement in levodopa-induced dyskinesia or decreased off episodes).
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

GROWTH HORMONES

Products Affected

- GENOTROPIN MINIQUICK SUBCUTANEOUS PREFILLED SYRINGE
- GENOTROPIN SUBCUTANEOUS CARTRIDGE
- HUMATROPE INJECTION CARTRIDGE
- NGENLA
- NORDITROPIN FLEXPRO SUBCUTANEOUS SOLUTION PEN-INJECTOR
- NUTROPIN AQ NUSPIN 10 SUBCUTANEOUS SOLUTION PEN-INJECTOR
- NUTROPIN AQ NUSPIN 20 SUBCUTANEOUS SOLUTION PEN-INJECTOR
- NUTROPIN AQ NUSPIN 5 SUBCUTANEOUS SOLUTION PEN-INJECTOR
- OMNITROPE SUBCUTANEOUS SOLUTION CARTRIDGE
- OMNITROPE SUBCUTANEOUS SOLUTION RECONSTITUTED
- SKYTROFA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	For new starts for growth hormone deficiency: Documentation showing bone age testing, height, weight, and Growth Hormone Stimulation Test results OR Insulin Growth Factor 1 level. For continuation of therapy or reauthorization for growth hormone deficiency: documentation (medical records) showing positive response to treatment.
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be an endocrinologist or nephrologist.
Coverage Duration	New starts will be authorized for 6 months. Cont of therapy or reauth until end of contract year.
Other Criteria	For new starts for growth hormone deficiency: 1) If the request is not for Genotropin, trial of, contraindication to, or medical reason for not using Genotropin. For requests for all other medically accepted indications other than growth hormone deficiency, the request will be approved for products other than Skytrofa.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

Formulary ID 25406

Last Update: 10/17/2025

PA Criteria	Criteria Details
Part B Prerequisite	No

HADLIMA

Products Affected

- HADLIMA
- HADLIMA PUSHTOUCH

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Specialist for submitted diagnosis.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	For ankylosing spondylitis: Trial of, medical reason for not using, or contraindication to naproxen. For Crohns Disease: Trial of, medical reason for not using (i.e. severe Crohns disease), or contraindication to 1 of the following: mercaptopurine, azathioprine, sulfasalazine, methotrexate or corticosteroid (e.g., prednisone, methylprednisolone). For pJIA: Trial of, medical reason for not using, or contraindication to 1 of the following DMARDs: methotrexate or leflunomide. For RA: Trial of, medical reason for not using, or contraindication to 1 disease modifying antirheumatic drug (DMARD) (methotrexate, leflunomide, or sulfasalazine). For UC: Trial of, medical reason for not using, or contraindication to 1 of the following conventional therapies: mercaptopurine, an aminosalicylate (i.e. mesalamine, sulfasalazine, azathioprine), or a corticosteroid (i.e. prednisone, methylprednisolone). For PsA, psoriasis, Hidradenitis Suppurativa, or Uveitis: approve.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

HEREDITARY ANGIOEDEMA AGENTS

Products Affected

- CINRYZE
- HAEGARDA
- ORLADEYO

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be an allergist, immunologist, rheumatologist or hematologist.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	For continuation of therapy or reauthorization: Documentation has been provided that patient has clinically benefited from medication.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

HETLIOZ

Products Affected

- HETLIOZ LQ
- *tasimelteon*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	For new starts of non-24 hour sleep-wake cycle: 1) Member is totally blind with no perception of light, 2) diagnosis of non-24 confirmed by a physiologic circadian phase marker (ex: dim light melatonin onset, assessment of core body temp or measurement of urinary melatonin levels) OR actigraphy with evaluation of sleep logs. For continuation of therapy or reauthorization: documentation of clinical benefit from use of the drug. For night-time sleep disturbances in Smith-Magenis Syndrome (SMS): approve
Age Restrictions	N/A
Prescriber Restrictions	Provider is a sleep specialist or neurologist.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

HIGH RISK MEDICATION

Products Affected

- *benztropine mesylate oral*
- *cyproheptadine hcl oral*
- *diphenoxylate-atropine oral liquid*
- *diphenoxylate-atropine oral tablet 2.5-0.025 mg*
- *dipyridamole oral*
- *ergotamine-caffeine*
- *glyburide micronized oral tablet 1.5 mg, 3 mg, 6 mg*
- *glyburide oral tablet 1.25 mg, 2.5 mg, 5 mg*
- *glyburide-metformin oral tablet 1.25-250 mg, 2.5-500 mg*
- *glyburide-metformin oral tablet 5-500 mg*
- *guanfacine hcl er*
- *guanfacine hcl oral*
- *hydroxyzine hcl oral syrup*
- *hydroxyzine hcl oral tablet 25 mg, 50 mg*
- *hydroxyzine pamoate oral*
- *indomethacin oral capsule 25 mg, 50 mg*
- *ketorolac tromethamine oral*
- *megestrol acetate oral suspension*
- *nifedipine oral*
- *promethazine hcl oral solution 6.25 mg/5ml*
- *promethazine hcl oral syrup*
- *promethazine hcl oral tablet*
- *promethazine hcl rectal suppository 12.5 mg, 25 mg*
- *promethazine-phenylephrine*
- *promethegan rectal suppository 50 mg*
- *trihexyphenidyl hcl*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	For patients 65 years old and older the prescriber has documented: 1) the benefits of treatment with the drug outweigh the potential risks identified for people 65 years old and older, and 2) the risks and side effects have been discussed and will be monitored.
Age Restrictions	Prior authorization only applies to members 65 years old or older.
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

PA Criteria	Criteria Details
Part B Prerequisite	No

HIGH RISK MEDICATION - PROTECTED CLASS DRUGS

Products Affected

- *estradiol oral*
- *estradiol transdermal patch twice weekly*
- *estradiol transdermal patch weekly*
- *megestrol acetate oral tablet*
- MENEST
- *perphenazine-amitriptyline*
- *phenobarbital oral elixir 20 mg/5ml*
- *phenobarbital oral tablet*
- PREMARIN ORAL

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	For patients 65 years old and older the prescriber has documented: 1) the benefits of treatment with the drug outweigh the potential risks identified for people 65 years old and older, and 2) the risks and side effects have been discussed and will be monitored.
Age Restrictions	Prior authorization only applies to members 65 years old or older.
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

HIGH RISK MEDICATION, BUTALBITAL

Products Affected

- *bac (butalbital-acetamin-caff)*
- *butalbital-acetaminophen oral tablet 50-325 mg*
- *butalbital-apap-caff-cod oral capsule 50-325-40-30 mg*
- *butalbital-apap-caffeine oral capsule 50-325-40 mg*
- *butalbital-apap-caffeine oral solution*
- *butalbital-apap-caffeine oral tablet 50-325-40 mg*
- *butalbital-asa-caff-codeine*
- *butalbital-aspirin-caffeine oral capsule*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	For patients 65 years old and older the prescriber has documented: 1) the benefits of treatment with the drug outweigh the potential risks identified for people 65 years old and older, and 2) the risks and side effects have been discussed and will be monitored.
Age Restrictions	Prior authorization only applies to members 65 years old or older.
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	Trial of, contraindication to, or medical reason for not using an oral NSAID.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

HIGH RISK MEDICATION, SHORT TERM MUSCLE RELAXANT

Products Affected

- *carisoprodol oral*
- *chlorzoxazone oral tablet 500 mg*
- *cyclobenzaprine hcl oral tablet 10 mg, 5 mg*
- *metaxalone oral tablet 800 mg*
- *methocarbamol oral tablet 500 mg, 750 mg*
- *orphenadrine citrate er*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	For patients 65 years old and older the prescriber has documented: 1) the benefits of treatment with the drug outweigh the potential risks identified for people 65 years old and older, and 2) the risks and side effects have been discussed and will be monitored.
Age Restrictions	Prior authorization only applies to members 65 years old or older.
Prescriber Restrictions	N/A
Coverage Duration	New starts will be authorized for 30 days. Continuation of therapy or reauth will be for 90 days.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

HIGH RISK MEDICATION, SLEEP AGENTS

Products Affected

- *eszopiclone*
- *temazepam*
- *zaleplon*
- *zolpidem tartrate er*
- *zolpidem tartrate oral tablet 10 mg*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	For patients 65 years old and older the prescriber has documented: 1) the benefits of treatment with the drug outweigh the potential risks identified for people 65 years old and older, and 2) the risks and side effects have been discussed and will be monitored. For zolpidem immediate release 10mg and zolpidem ER: trial of or medical reason for not using zolpidem immediate release 5mg.
Age Restrictions	Prior authorization only applies to members 65 years old or older.
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

HUMIRA

Products Affected

- HUMIRA (1 PEN)
- HUMIRA (2 PEN) SUBCUTANEOUS AUTO-INJECTOR KIT
- HUMIRA (2 SYRINGE) SUBCUTANEOUS PREFILLED SYRINGE KIT 10 MG/0.1ML, 20 MG/0.2ML, 40 MG/0.4ML, 40 MG/0.8ML
- HUMIRA-CD/UC/HS STARTER SUBCUTANEOUS AUTO-INJECTOR KIT 80 MG/0.8ML
- HUMIRA-PED>=40KG UC STARTER SUBCUTANEOUS AUTO-INJECTOR KIT
- HUMIRA-PSORIASIS/UEVIT STARTER SUBCUTANEOUS AUTO-INJECTOR KIT

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Specialist for submitted diagnosis.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	For ankylosing spondylitis: Trial of, medical reason for not using, or contraindication to naproxen. For Crohns Disease: Trial of, medical reason for not using (i.e. severe Crohns disease), or contraindication to 1 of the following: mercaptopurine, azathioprine, sulfasalazine, methotrexate or corticosteroid (e.g., prednisone, methylprednisolone). For pJIA: Trial of, medical reason for not using, or contraindication to 1 of the following DMARDs: methotrexate or leflunomide. For RA: Trial of, medical reason for not using, or contraindication to 1 disease modifying antirheumatic drug (DMARD) (methotrexate, leflunomide, or sulfasalazine). For UC: Trial of, medical reason for not using, or contraindication to 1 of the following conventional therapies: mercaptopurine, an aminosalicilate (i.e. mesalamine, sulfasalazine, azathioprine), or a corticosteroid (i.e. prednisone, methylprednisolone). For PsA, psoriasis, Hidradenitis Suppurativa, or Uveitis: approve.

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

HYFTOR

Products Affected

- HYFTOR

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	For new starts: documentation of diagnosis of tuberous sclerosis with facial angiofibroma. For continuation of therapy or reauthorization: documentation that the member has experienced a clinical benefit from treatment (e.g. improvement in size and color of angiofibroma).
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a neurologist or provider who specializes in the treatment of genetic or dermatologic disorders.
Coverage Duration	New starts: 3 months. Cont. of therapy or reauthorization: until end of contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

ICATIBANT

Products Affected

- *icatibant acetate subcutaneous solution prefilled syringe*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be an immunologist, allergist, rheumatologist, or hematologist.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

ILARIS

Products Affected

- ILARIS SUBCUTANEOUS SOLUTION

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Documentation was submitted indicating that the member was evaluated for active or latent TB infection (i.e. tuberculin skin test)
Age Restrictions	N/A
Prescriber Restrictions	Specialist for submitted diagnosis.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	For sJIA: approve.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

ILUMYA

Products Affected

- ILUMYA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Specialist for submitted diagnosis.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	For psoriasis: Either 1) Trial of, medical reason for not using, or contraindication to 2 of the following therapies: Stelara, Skyrizi, Tremfya, Enbrel, Hadlima, or Humira, or 2) If utilized within the past 120 days, approve for continuation of therapy.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

IMBRUVICA

Products Affected

- IMBRUVICA ORAL CAPSULE
- IMBRUVICA ORAL TABLET 140 MG, 280 MG, 420 MG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be an oncologist or specialist for submitted diagnosis.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	For new starts for treatment of graft-versus-host disease (GVHD): Trial of, contraindication to, or medical reason for not using a systemic corticosteroid. For continuation of therapy of for treatment of GVHD: documentation of clinical benefit from use of the drug (i.e. symptom improvement, reduction in corticosteroid dose). For all other indications, approve.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

IMPAVIDO

Products Affected

- IMPAVIDO

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Documentation of diagnosis with one of the following: (a) Visceral leishmaniasis due to <i>Leishmania donovani</i> , (b) Cutaneous leishmaniasis due to <i>Leishmania braziliensis</i> , <i>Leishmania guyanensis</i> , or <i>Leishmania panamensis</i> , (c) Mucosal leishmaniasis due to <i>Leishmania braziliensis</i> .
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized for 28 days.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

INCRELEX

Products Affected

- INCRELEX

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an endocrinologist.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

JAKAFI

Products Affected

- JAKAFI

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be an oncologist or specialist for submitted diagnosis.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	For new starts for treatment of graft-versus-host disease (GVHD): Trial of, contraindication to, or medical reason for not using a systemic corticosteroid. For continuation of therapy of for treatment of GVHD: documentation of clinical benefit from use of the drug (i.e. symptom improvement, reduction in corticosteroid dose). For all other indications, approve.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

JYLAMVO

Products Affected

- JYLAMVO

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be an oncologist, a rheumatologist, a dermatologist, or other appropriate specialist
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

KALYDECO

Products Affected

- KALYDECO

PA Criteria	Criteria Details
Exclusion Criteria	Combination use with Orkambi, Symdeko, or Trikafta.
Required Medical Information	Documentation of CFTR gene that is responsive to ivacaftor treatment.
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be a pulmonologist or an expert in the treatment of cystic fibrosis.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

KERENDIA

Products Affected

- KERENDIA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	New starts will be authorized for 3 months. Cont of therapy or reauth until end of contract year.
Other Criteria	For new starts: 1) Documentation of diagnosis of chronic kidney disease due to type 2 diabetes mellitus AND 2) Documentation of serum potassium levels less than or equal to 5 mEq/L AND 3) eGFR greater than or equal to 25ml/min/1.73 m2 AND 4) Documentation that member is taking Kerendia in combination with an angiotensin-converting enzyme inhibitor (ACEi) or angiotensin receptor blocker (ARB) at maximum tolerated doses or documentation has been provided that the member is unable to tolerate ACEi or ARB AND 4) Documented trial of, contraindication to, or medical reason for not using a sodium-glucose cotransporter-2 (SGLT2) inhibitor. For continuation of therapy or reauthorization: 1) Documentation of serum potassium levels less than or equal to 5.5 mEq/L AND 2) Documentation that member is taking Kerendia in combination with an ACEi or ARB at maximum tolerated doses or documentation has been provided that the member is unable to tolerate ACEi or ARB.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

KEVZARA

Products Affected

- KEVZARA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Specialist for submitted diagnosis.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	For RA: Either 1) Trial of, medical reason for not using, or contraindication to 2 of the following: Enbrel, Humira, Hadlima, Rinvoq or Xeljanz or 2) If utilized within the past 120 days, approve for continuation of therapy. For polymyalgia rheumatica (PMR): Trial of, medical reason for not using, or contraindication to corticosteroids. For pJIA: Either 1) Trial of, medical reason for not using, or contraindication to 2 of the following: Enbrel, Humira, Hadlima, or Xeljanz or 2) If utilized within the past 120 days, approve for continuation of therapy.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

KINERET

Products Affected

- KINERET SUBCUTANEOUS
SOLUTION PREFILLED SYRINGE

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Specialist for submitted diagnosis.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	For RA: Either 1) Trial of, medical reason for not using, or contraindication to 2 of the following: Enbrel, Humira, Hadlima, Rinvoq or Xeljanz or 2) If utilized within the past 120 days, approve for continuation of therapy. For cryopyrin-associated periodic syndromes or deficiency of interleukin-1 receptor antagonist: Approve.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

LEQSELVI

Products Affected

- LEQSELVI

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Documentation was submitted indicating that the member was evaluated for active or latent TB infection (i.e. tuberculin skin test).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	Documentation of confirmed diagnosis and other causes of hair loss have been ruled out.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

LIBERVANT

Products Affected

- LIBERVANT

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	Patient is between 2 to 5 years of age.
Prescriber Restrictions	Prescriber must be a neurologist.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

LITFULO

Products Affected

- LITFULO

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Documentation was submitted indicating that the member was evaluated for active or latent TB infection (i.e. tuberculin skin test)
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	Documentation of confirmed diagnosis and other causes of hair loss have been ruled out.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

LIVMARLI

Products Affected

- LIVMARLI

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a gastroenterologist or hepatologist.
Coverage Duration	New starts will be authorized for 6 months. Cont of therapy or reauth until end of contract year.
Other Criteria	For new starts: 1) Trial of, contraindication to, or medical reason for not using both of the following: cholestyramine AND rifampin. 2) Prescriber attests that the member has cholestasis 3) Baseline serum bile acid level is provided. 4) Documentation of patients weight. For continuation of therapy or reauthorization: 1) Documentation submitted indicating the member has had all of the following: an improvement in pruritis (e.g. improved observed scratching, decreased sleep disturbances/nighttime awakenings due to scratching, etc.) AND reduction in serum bile acid level from baseline. 2) Prescriber attests that patient has had no evidence of hepatic decompensation (e.g. variceal hemorrhage, ascites, hepatic encephalopathy, portal hypertension, etc.). 3) Documentation of patients weight.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

LIVTENSITY

Products Affected

- LIVTENCITY

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Cytomegalovirus (CMV): (1) Documented diagnosis of CMV infection AND (2) Member is a recipient of one of the following: (a) hematopoietic stem cell transplant, (b) solid organ transplant AND (3) patient has tried and failed treatment with valganciclovir, ganciclovir, cidofovir, or foscarnet.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a transplant, oncologist, or infectious disease specialist.
Coverage Duration	Request will be authorized for 8 weeks.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

LODOCO

Products Affected

- LODOCO

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be, or in consultation with a specialist in the treatment of cardiovascular disease, such as a cardiologist
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	Documentation that patient has established atherosclerotic disease or multiple risk factors for cardiovascular disease AND documentation that patient does not have pre-existing blood dyscrasias (ex. leukopenia, thrombocytopenia) and patient does not have renal failure (CrCl less than 15 ml/min) or severe hepatic impairment
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

LUCEMYRA

Products Affected

- *lofexidine hcl*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized for 14 days.
Other Criteria	Patient must have trial of, contraindication to, or medical reason for not using clonidine.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

LUPKYNIS

Products Affected

- LUPKYNIS

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent use with cyclophosphamide.
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be rheumatologist, nephrologist, or other specialist in the treatment of autoimmune disorders.
Coverage Duration	New starts will be authorized for 6 months. Cont of therapy or reauth until end of contract year.
Other Criteria	For new starts: 1) Documentation of urine protein/creatinine ratio (UPCR), 2) Documentation that the member has a baseline eGFR greater than 45 mL/min/1.73m ² or that benefit outweighs risk of using this medication at current eGFR, and 3) Concurrent use of or medical reason for not using background immunosuppressive therapy regimen. For continuation of therapy or reauthorization: Documentation of improvement in renal function (i.e. reduction in UPCR or no confirmed decrease from baseline eGFR greater than or equal to 20%).
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

LYBALVI

Products Affected

- LYBALVI

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with opioids.
Required Medical Information	Attestation from the provider that the member has had an opioid-free period of a minimum of 7 days after last use of shorting-acting opioids and 14 days from last use of long-acting opioids before initiating Lybalvi.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	Documented trial of, contraindication to, or medical reason for not using at least two generic antipsychotics, one of which must be generic olanzapine.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

MAVYRET

Products Affected

- MAVYRET

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Detectable HCV RNA viral load prior to treatment within 6 months of request. In addition, documentation of treatment history, and if cirrhotic, documentation of compensated or decompensated cirrhosis.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized for 8-16 weeks as per AASLD-IDSA guidance.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

METHYLTESTOSTERONE

Products Affected

- *methyltestosterone oral*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

METYROSIONE

Products Affected

- *metyrosine*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Documentation of one of the following: 1) Concurrent use of alpha adrenergic blockers, 2) Medical reason for being unable to use an alpha adrenergic blocker, OR 3) Patient is not a candidate for surgical resection and requires long term treatment with metyrosine.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

MIFEPRISTONE

Products Affected

- *mifepristone oral tablet 300 mg*

PA Criteria	Criteria Details
Exclusion Criteria	For all members patient must not be currently on simvastatin, lovastatin, cyclosporine, dihydroergotamine, ergotamine, fentanyl, pimozide, quinidine, sirolimus, and tacrolimus.
Required Medical Information	Reviewer will verify available claim history to confirm member is not taking simvastatin, lovastatin, cyclosporine, dihydroergotamine, ergotamine, fentanyl, pimozide, quinidine, sirolimus or tacrolimus concurrently with mifepristone.
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be an endocrinologist.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

MIGLUSTAT

Products Affected

- *miglustat*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	For new starts, documentation of diagnosis for mild to moderate type 1 Gaucher disease. For continuation of therapy or reauthorization: documentation of clinical benefit from use of the drug (i.e. increased platelet count, improvement in anemia, PFT's, improvement in radiographic scans, improved quality of life).
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a specialist in treatment of Gaucher's disease
Coverage Duration	New starts will be authorized for 6 months. Cont of therapy or reauth until end of contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

MULTIPLE SCLEROSIS AGENTS

Products Affected

- BAFIERTAM
- BETASERON SUBCUTANEOUS KIT
- *dimethyl fumarate oral capsule delayed release 120 mg, 240 mg*
- *dimethyl fumarate starter pack oral capsule delayed release therapy pack*
- *fingolimod hcl*
- *glatiramer acetate subcutaneous solution prefilled syringe 20 mg/ml, 40 mg/ml*
- *glatopa subcutaneous solution prefilled syringe 20 mg/ml, 40 mg/ml*
- KESIMPTA
- MAVENCLAD (10 TABS)
- MAVENCLAD (4 TABS)
- MAVENCLAD (5 TABS)
- MAVENCLAD (6 TABS)
- MAVENCLAD (7 TABS)
- MAVENCLAD (8 TABS)
- MAVENCLAD (9 TABS)
- MAYZENT
- MAYZENT STARTER PACK
- PONVORY
- PONVORY STARTER PACK
- REBIF REBIDOSE SUBCUTANEOUS SOLUTION AUTO-INJECTOR
- REBIF REBIDOSE TITRATION PACK SUBCUTANEOUS SOLUTION AUTO-INJECTOR
- REBIF SUBCUTANEOUS SOLUTION PREFILLED SYRINGE
- REBIF TITRATION PACK SUBCUTANEOUS SOLUTION PREFILLED SYRINGE
- TASCENSO ODT
- *teriflunomide*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Specialist for submitted diagnosis.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	If the medication request is for glatiramer, Glatopa, or dimethyl fumarate, the request will be approved. If the member is over 17 years of age and the request is not for glatiramer, Glatopa, or dimethyl fumarate for multiple sclerosis, the member must have a documented trial of, contraindication to or a medical reason for not using 2 of the following dimethyl fumarate, glatiramer, Glatopa, teriflunomide, or fingolimod. If the request is for

PA Criteria	Criteria Details
	fingolimod and the member is 17 years of age or younger, the request will be approved.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

MYFEMBREE

Products Affected

- MYFEMBREE

PA Criteria	Criteria Details
Exclusion Criteria	Patient has history of osteoporosis or hepatic impairment.
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be an OB, gynecologist or reproductive endocrinologist.
Coverage Duration	New starts will be authorized for 6 months. Cont of therapy or reauth until end of contract year.
Other Criteria	For new starts for menorrhagia: Trial of, contraindication to, or medical reason for not using an estrogen-progestin contraceptive therapy. For new starts if one of the following drugs has been tried previously, a trial of estrogen-progestin contraceptive therapy is not required: gonadotropin-releasing hormone (GnRH) agonists or tranexamic acid. New starts for endometriosis: Trial of, contraindication to, or medical reason for not using the following concurrently for endometriosis: analgesic pain reliever (e.g. NSAIDs, COX-2 inhibitors) AND either combined estrogen-progestin oral contraceptive, progestin (e.g. medroxyprogesterone acetate, norethindrone), gonadotropin-releasing hormone (GnRH) agonists (e.g. Lupron Depot), OR danazol. For continuation of therapy or reauthorization both of the following are required: 1) Treatment does not exceed the eligible maximum lifetime treatment duration of 2 years, and 2) Documentation has been provided that the member has obtained clinical benefit from medication (e.g. reduced menstrual bleeding from baseline, pain relief).
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

NASAL ANTISEIZURE AGENTS

Products Affected

- NAYZILAM
- VALTOCO 10 MG DOSE
- VALTOCO 15 MG DOSE NASAL LIQUID THERAPY PACK 2 X 7.5 MG/0.1ML
- VALTOCO 20 MG DOSE NASAL LIQUID THERAPY PACK 2 X 10 MG/0.1ML
- VALTOCO 5 MG DOSE

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a neurologist.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

NITISINONE

Products Affected

- *nitisinone*
- ORFADIN ORAL SUSPENSION

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be a geneticist, metabolic specialist, hepatologist, or liver transplant specialist.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

NON-AMPHETAMINE CENTRAL NERVOUS SYSTEM AGENTS

Products Affected

- *armodafinil*
- *modafinil oral*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

NUCALA

Products Affected

- NUCALA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Specialist for submitted diagnosis.
Coverage Duration	New starts will be authorized for 6 months. Cont of therapy or reauth until end of contract year.
Other Criteria	New starts for severe asthma: 1) Baseline blood eosinophil count greater than or equal to 150 cells per microliter AND 2) symptoms with equal to or greater than 1 exacerbations in the previous 12 months requiring additional medical treatment, (e.g. oral systemic steroids) while on a high-dose inhaled corticosteroid with an additional controller medication (ie. long-acting B2 agonist). New starts for eosinophilic granulomatosis with polyangiitis (EGPA): trial of, contraindication to, or medical reason for not using one of the following medications: cyclophosphamide or methotrexate. New starts for hypereosinophilic syndrome without an identifiable non-hematologic secondary cause: 1) 2 or more flares within the past 12 months AND 2) trial of, contraindication to, or medical reason for not using oral corticosteroids. New starts for chronic rhinosinusitis with nasal polyps: trial of, contraindication to, or medical reason for not using nasal corticosteroids OR member has had prior surgery for nasal polyps. Continuation of therapy or re-authorization for all indications: clinical benefit from use of the drug.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID 25406

Last Update: 10/17/2025

NUEDEXTA

Products Affected

- NUEDEXTA

PA Criteria	Criteria Details
Exclusion Criteria	Complete atrioventricular (AV) block without implanted pacemaker, or at high risk of complete AV block. History of heart failure. Concomitant use with MAOIs or use of MAOIs within 14 days. Concomitant use with drugs containing quinidine, quinine, or mefloquine. History of quinine-, mefloquine-, dextromethorphan/quinidine-, or quinidine-induced thrombocytopenia, hepatitis, bone marrow depression, or lupus-like syndrome. Non-Part D indications.
Required Medical Information	Confirmation diagnosis is for Part D indication.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a neurologist or psychiatrist.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

NUPLAZID

Products Affected

- NUPLAZID ORAL CAPSULE
- NUPLAZID ORAL TABLET 10 MG

PA Criteria	Criteria Details
Exclusion Criteria	Patient has a history of dementia-related psychosis.
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

OCREVUS

Products Affected

- OCREVUS ZUNOVO

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	18 years of age and older.
Prescriber Restrictions	Specialist for submitted diagnosis.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	For new starts for Clinically Isolated Syndrome (CIS), Relapsing Remitting Multiple Sclerosis (RRMS), or Secondary Progressive Multiple Sclerosis (SPMS): 1) Documentation of CIS, RRMS, or SPMS AND 2) The member must have a documented trial of, contraindication to, or medical reason for not using both dimethyl fumarate AND glatiramer or Glatopa. For new starts for Primary Progressive Multiple Sclerosis (PPMS): Documentation of PPMS. For all continuation of therapy or reauthorization: Documentation that the prescriber has evaluated the member and recommends continuation of therapy (clinical benefit).
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

OCTREOTIDE

Products Affected

- *octreotide acetate injection solution 100 mcg/ml, 1000 mcg/ml, 200 mcg/ml, 50 mcg/ml, 500 mcg/ml*
- *octreotide acetate intramuscular*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	For new starts for acromegaly: pt meets one of the following (1) inadequate response to surgery and/or radiotherapy OR (2) pt is not an appropriate candidate for surgery and/or radiotherapy OR (3) pt is experiencing negative effects due to tumor size (ex: optic nerve compression). Continuation of therapy or reauthorization: documentation of clinical improvement with therapy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	Continuation of therapy or reauthorization: documentation of clinical improvement with therapy.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

OFEV

Products Affected

- OFEV

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be a pulmonologist or lung transplant specialist.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	For a diagnosis of idiopathic pulmonary fibrosis: 1) Documentation of disease as demonstrated on a high resolution CT scan or through lung biopsy and 2) Documented trial of, contraindication to, or medical reason for not using pirfenidone. For a diagnosis of systemic sclerosis-associated interstitial lung disease (SSc-ILD): documented trial of, contraindication to, or medical reason for not using mycophenolate mofetil or cyclophosphamide. For a diagnosis of chronic fibrosing interstitial lung diseases (ILDs) with a progressive phenotype: documentation is provided confirming diagnosis.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

OPDIVO QVANTIG

Products Affected

- OPDIVO QVANTIG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be an oncologist or specialist for submitted diagnosis.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

OPSUMIT

Products Affected

- OPSUMIT

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Documentation of pulmonary arterial hypertension (PAH) WHO Group I and PAH Functional Class.
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be a pulmonologist or cardiologist.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	Trial of, contraindication to, or medical reason for not using sildenafil.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

ORAL ANTINEOPLASTIC AGENTS

Products Affected

- *abiraterone acetate*
- ABIRTEGA
- AKEEGA
- ALECENSA
- ALUNBRIG
- AUGTYRO
- AVMAPKI FAKZYNJA CO-PACK
- AYVAKIT
- BALVERSA
- *bexarotene*
- BOSULIF
- BRAFTOVI ORAL CAPSULE 75 MG
- BRUKINSA
- CABOMETYX
- CALQUENCE ORAL TABLET
- CAPRELSA
- COMETRIQ (100 MG DAILY DOSE) ORAL KIT 80 & 20 MG
- COMETRIQ (140 MG DAILY DOSE) ORAL KIT 3 X 20 MG & 80 MG
- COMETRIQ (60 MG DAILY DOSE)
- COPIKTRA
- COTELLIC
- DANZITEN
- *dasatinib*
- DAURISMO
- ERIVEDGE
- ERLEADA
- *erlotinib hcl*
- EULEXIN
- *everolimus oral tablet 10 mg, 2.5 mg, 5 mg, 7.5 mg*
- *everolimus oral tablet soluble*
- FOTIVDA
- FRUZAQLA
- GAVRETO
- *gefitinib*
- GILOTRIF
- GOMEKLI
- HERNEXEOS
- IBRANCE
- IBTROZI
- ICLUSIG
- IDHIFA
- *imatinib mesylate oral*
- IMBRUVICA ORAL SUSPENSION
- IMKELDI
- INLYTA
- INQOVI
- INREBIC
- ITOVEBI
- IWILFIN
- JAYPIRCA
- KISQALI (200 MG DOSE)
- KISQALI (400 MG DOSE)
- KISQALI (600 MG DOSE)
- KISQALI FEMARA (200 MG DOSE)
- KISQALI FEMARA (400 MG DOSE)
- KISQALI FEMARA (600 MG DOSE)
- KOSELUGO ORAL CAPSULE
- KRAZATI
- *lapatinib ditosylate*
- LAZCLUZE
- *lenalidomide*
- LENVIMA (10 MG DAILY DOSE)
- LENVIMA (12 MG DAILY DOSE)
- LENVIMA (14 MG DAILY DOSE)
- LENVIMA (18 MG DAILY DOSE)
- LENVIMA (20 MG DAILY DOSE)
- LENVIMA (24 MG DAILY DOSE)
- LENVIMA (4 MG DAILY DOSE)
- LENVIMA (8 MG DAILY DOSE)
- LEUKERAN
- LONSURF
- LORBRENA
- LUMAKRAS
- LYNPARZA ORAL TABLET
- LYTGobi (12 MG DAILY DOSE)
- LYTGobi (16 MG DAILY DOSE)
- LYTGobi (20 MG DAILY DOSE)
- MEKINIST
- MEKTOVI
- *mercaptopurine oral suspension*
- MODEYSO

Formulary ID 25406

Last Update: 10/17/2025

- NERLYNX
- *nilotinib d-tartrate*
- *nilotinib hcl*
- *nilutamide*
- NINLARO
- NUBEQA
- ODOMZO
- OGSIVEO
- OJEMDA
- OJJAARA
- ONUREG
- ORGOVYX
- ORSERDU
- *pazopanib hcl*
- PEMAZYRE
- PIQRAY (200 MG DAILY DOSE)
- PIQRAY (250 MG DAILY DOSE)
- PIQRAY (300 MG DAILY DOSE)
- POMALYST
- QINLOCK
- RETEVMO
- REVLIMID
- REVUFORJ
- REZLIDHIA
- ROMVIMZA
- ROZLYTREK
- RUBRACA
- RYDAPT
- SCEMBLIX
- SOLTAMOX
- *sorafenib tosylate*
- STIVARGA
- *sunitinib malate*
- TABLOID
- TABRECTA
- TAFINLAR
- TAGRISSO
- TALZENNA
- TAZVERIK
- TEPMETKO
- THALOMID
- TIBSOVO
- *toremifene citrate*
- *tretinoin oral*
- TRUQAP
- TUKYSA
- TURALIO ORAL CAPSULE 125 MG
- VANFLYTA
- VENCLEXTA
- VENCLEXTA STARTING PACK
- VERZENIO
- VITRAKVI
- VIZIMPRO
- VONJO
- VORANIGO
- WELIREG
- XALKORI
- XOSPATA
- XPOVIO (100 MG ONCE WEEKLY)
ORAL TABLET THERAPY PACK 50
MG
- XPOVIO (40 MG ONCE WEEKLY)
ORAL TABLET THERAPY PACK 10
MG, 40 MG
- XPOVIO (40 MG TWICE WEEKLY)
ORAL TABLET THERAPY PACK 40
MG
- XPOVIO (60 MG ONCE WEEKLY)
ORAL TABLET THERAPY PACK 60
MG
- XPOVIO (60 MG TWICE WEEKLY)
- XPOVIO (80 MG ONCE WEEKLY)
ORAL TABLET THERAPY PACK 40
MG
- XPOVIO (80 MG TWICE WEEKLY)
- XTANDI
- YONSA
- ZEJULA ORAL TABLET
- ZELBORAF
- ZOLINZA
- ZYDELIG
- ZYKADIA ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be an oncologist or specialist for submitted diagnosis.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

ORAL ANTIPSYCHOTICS

Products Affected

- CAPLYTA
- COBENFY
- COBENFY STARTER PACK
- FANAPT
- FANAPT TITRATION PACK A
- FANAPT TITRATION PACK B ORAL TABLET
- FANAPT TITRATION PACK C ORAL TABLET
- OPIPZA ORAL FILM 10 MG, 2 MG, 5 MG
- VRAYLAR ORAL CAPSULE

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	For schizophrenia, manic or mixed episodes associated with bipolar I disorder, major depressive disorder associated with bipolar I or II disorder, adjunctive treatment of major depressive disorder, irritability associated with autistic disorder or treatment of Tourette's disorder: trial of, contraindication to, or medical reason for not using two generic antipsychotics. If the request is for Vraylar for major depressive disorder: provider attestation that the member is concurrently using an antidepressant.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

ORENCIA

Products Affected

- ORENCIA CLICKJECT
- ORENCIA INTRAVENOUS
- ORENCIA SUBCUTANEOUS SOLUTION PREFILLED SYRINGE

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Specialist for submitted diagnosis.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	For pJIA: Either 1) Trial of, medical reason for not using, or contraindication to 2 of the following: Enbrel, Humira, Hadlima, or Xeljanz or 2) If utilized within the past 120 days, approve for continuation of therapy. For PsA: Either 1) Trial of, medical reason for not using, or contraindication to 2 of the following therapies: Stelara, Skyrizi, Tremfya, Xeljanz, Rinvoq, Enbrel, Hadlima, or Humira, or 2) If utilized within the past 120 days, approve for continuation of therapy. For RA: Either 1) Trial of, medical reason for not using, or contraindication to 2 of the following: Enbrel, Humira, Hadlima, Rinvoq or Xeljanz or 2) If utilized within the past 120 days, approve for continuation of therapy. For acute graft versus host disease: Attestation member is taking in combination with a calcineurin inhibitor and methotrexate.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

ORIAHNN

Products Affected

- ORIAHNN

PA Criteria	Criteria Details
Exclusion Criteria	Patient has history of osteoporosis or hepatic impairment.
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be an OB, gynecologist or reproductive endocrinologist.
Coverage Duration	New starts will be authorized for 6 months. Cont of therapy or reauth until end of contract year.
Other Criteria	For new starts: Trial of, contraindication to, or medical reason for not using an estrogen-progestin contraceptive therapy. For new starts if one of the following drugs has been tried previously, a trial of estrogen-progestin contraceptive therapy is not required: gonadotropin-releasing hormone (GnRH) agonists or tranexamic acid. For continuation of therapy or reauthorization both of the following are required: 1) Treatment does not exceed the eligible maximum lifetime treatment duration of 2 years, and 2) Documentation has been provided that the member has obtained clinical benefit from medication (e.g. reduced menstrual bleeding from baseline).
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

ORILISSA

Products Affected

- ORILISSA

PA Criteria	Criteria Details
Exclusion Criteria	Patient has osteoporosis or severe hepatic impairment.
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be an OB or gynecologist.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	Trial of, contraindication to, or medical reason for not using the following concurrently for endometriosis: analgesic pain reliever (e.g. NSAIDs, COX-2 inhibitors) AND either combined estrogen-progestin oral contraceptive, progestin (e.g. medroxyprogesterone acetate, norethindrone), gonadotropin-releasing hormone (GnRH) agonists (e.g. Lupron Depot), OR danazol. For continuation of therapy or reauthorization both of the following are required: 1) Treatment does not exceed the eligible maximum lifetime treatment duration of 2 years for 150mg tablet or 6 months for 200mg tablet, and 2) Documentation has been provided that the member has obtained clinical benefit from the medication.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

ORKAMBI

Products Affected

- ORKAMBI

PA Criteria	Criteria Details
Exclusion Criteria	Combination use with Kalydeco, Symdeko, or Trikafta.
Required Medical Information	Documentation of CFTR gene that is responsive to lumacaftor-ivacaftor treatment.
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be a pulmonologist or an expert in the treatment of cystic fibrosis.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

OTEZLA

Products Affected

- OTEZLA
- OTEZLA XR
- OTEZLA/OTEZLA XR INITIATION PK

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Specialist for submitted diagnosis.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	For moderate to severe psoriasis: Either 1) Trial of, medical reason for not using, or contraindication to 2 of the following therapies: Stelara, Skyrizi, Tremfya, Enbrel, Hadlima, or Humira, or 2) If utilized within the past 120 days, approve for continuation of therapy. For PsA: Either 1) Trial of, medical reason for not using, or contraindication to 2 of the following therapies: Stelara, Skyrizi Tremfya, Xeljanz, Rinvoq, Enbrel, Hadlima, or Humira, or 2) If utilized within the past 120 days, approve for continuation of therapy. For Behcet's Syndrome or mild psoriasis: Approve.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

OXERVATE

Products Affected

- OXERVATE

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be an ophthalmologist.
Coverage Duration	Request will be authorized for 8 weeks.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

OXYCODONE ER

Products Affected

- *oxycodone hcl er oral tablet er 12 hour abuse-deterrent 10 mg, 20 mg, 40 mg, 80 mg*
- OXYCONTIN ORAL TABLET ER 12 HOUR ABUSE-DETERRENT 10 MG, 15 MG, 20 MG, 30 MG, 40 MG, 60 MG, 80 MG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	Members being treated for active cancer diagnoses, sickle cell diagnoses, those in hospice care, or receiving palliative care will be excluded from the concurrent benzodiazepine and muscle relaxant therapy requirement. For new starts, ALL of the following are required: (1) Member has documented history of receiving an immediate-release opioid, (2) Member has a documented trial of, contraindication to, or medical reason for not using long-acting morphine sulfate, (3) If member is on concurrent benzodiazepines and/or muscle relaxant therapy, the prescriber attests that concurrent therapy is medically necessary, (4) Member is not being treated for substance abuse with buprenorphine-containing products. For continuing therapy, ALL of the following are required: (1) Member's pain has been assessed within the last 6 months, (2) Member has demonstrated clinical improvement in pain and function on current medication regimen, (3) If member is on concurrent benzodiazepines and/or muscle relaxant therapy, the prescriber attests that concurrent therapy is medically necessary, (4) Member is not being treated for substance abuse with buprenorphine-containing products.
Indications	All Medically-accepted Indications.

PA Criteria	Criteria Details
Off-Label Uses	N/A
Part B Prerequisite	No

PCSK9 INHIBITORS

Products Affected

- PRALUENT SUBCUTANEOUS SOLUTION AUTO-INJECTOR
- REPATHA
- REPATHA SURECLICK

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be a cardiologist, endocrinologist, or a specialist in treatment of lipid disorders.
Coverage Duration	New starts will be authorized for 4 months. Cont of therapy or reauth until end of contract year.
Other Criteria	For ALL diagnoses (including primary hyperlipidemia) for new starts, attestations of the following: 1) Two fasting lipid panel reports within the past 12 months with abnormal LDL cholesterol results (above 70mg/dL) after treatment for a minimum of 3 months with a high potency statin (atorvastatin and rosuvastatin) and ezetimibe, or a medical reason (contraindication or intolerance) has been provided as to why the patient is unable to use these therapies, and 2) If patient experiences statin intolerance, trial of statin re-challenge with maximally tolerated dose of statins with continued abnormal LDL cholesterol results (above 70mg/dL) or with attestation of return of side effects. For familial hypercholesterolemia (FH), attestation of one of the following: 1) genetic testing confirming FH diagnosis, 2) a clinical diagnosis of 'definite' FH using the Dutch Lipid Clinic Diagnostic criteria, OR Simon-Broome Diagnostic criteria, OR American Heart Association criteria. For ASCVD, additional attestation of history of acute coronary syndromes, history of MI, stable or unstable angina, coronary or other arterial revascularization, stroke, TIA, or peripheral arterial disease presumed to be of atherosclerotic origin. For ALL diagnoses for continuation of therapy or reauthorization: attestation of improvement in LDL from new start.
Indications	All Medically-accepted Indications.

PA Criteria	Criteria Details
Off-Label Uses	N/A
Part B Prerequisite	No

PEGINTERFERON

Products Affected

- PEGASYS SUBCUTANEOUS SOLUTION 180 MCG/ML
- PEGASYS SUBCUTANEOUS SOLUTION PREFILLED SYRINGE

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	For Hepatitis C: 1) Labs within 3 months of request: liver function tests and detectable HCV RNA viral load. 2) Documentation of genotype, treatment history, and if cirrhotic, documentation of compensated or decompensated cirrhosis. For Hepatitis B: 1) Labs within 3 months of request: ALT/AST, and 2) HBeAg status. For polycythemia vera, approve.
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be a gastroenterologist, hepatologist, infectious disease doctor or transplant specialist.
Coverage Duration	Request will be authorized for 24 to 48 weeks as defined by compendia.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

PENICILLAMINE

Products Affected

- *penicillamine oral tablet*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Specialist for submitted diagnosis.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	For RA: Trial of, medical reason for not using, or contraindication to 2 of the following: Enbrel, Humira, Hadlima, Rinvoq or Xeljanz. For other indications, approve.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

PENTAMIDINE SOLUTION FOR INJECTION

Products Affected

- *pentamidine isethionate injection*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

PERSERIS

Products Affected

- PERSERIS

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	The member has a documented history of receiving oral risperidone without any clinically significant side effects.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	Trial of, contraindication to, or medical reason (e.g. intolerance, hypersensitivity or contraindication) for not using at least two of the following: Abilify Maintena, Abilify Asimtufii, Invega Sustenna, Invega Trinza, or Risperdion Microsphere ER.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

PHENOXYBENZAMINE

Products Affected

- *phenoxybenzamine hcl oral*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	Trial of, contraindication to, or medical reason for not using doxazosin.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

PIASKY

Products Affected

- PIASKY

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be a hematologist or specialist for submitted diagnosis.
Coverage Duration	New starts will be authorized for 6 months. Cont of therapy or reauth until end of contract year.
Other Criteria	For new starts: The member has a documented diagnosis of paroxysmal nocturnal hemoglobinuria (PNH) AND documentation of glycosylphosphatidylinositol-anchored proteins (GPI-APs) deficiency as demonstrated through flow cytometry. For continuation of therapy: Documentation that member has had positive response to therapy (e.g., improvement in hemoglobin levels, normalization of lactase dehydrogenase [LDH] levels).
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

PIRFENIDONE

Products Affected

- *pirfenidone*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	For idiopathic pulmonary fibrosis, documentaton of all of the following: 1) confirmation of diagnosis on high resolution CT scan or through lung biopsy AND 2) FVC greater than or equal to 50% of the predicted value.
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be a pulmonologist or lung transplant specialist.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

POSACONAZOLE

Products Affected

- *posaconazole oral*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Documentation of a consultation with an infectious disease specialist, a transplant specialist, or an oncologist.
Coverage Duration	28 days for oropharyngeal candidiasis, end of contract year for other indications
Other Criteria	For treatment of oropharyngeal candidiasis: trial of, contraindication to, or medical reason for not using fluconazole or itraconazole. For prophylaxis of invasive aspergillus infections due to being severely immunocompromised: trial of, contraindication to, or medical reason for not using voriconazole.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

PRETOMANID

Products Affected

- PRETOMANID

PA Criteria	Criteria Details
Exclusion Criteria	MDR-TB that is not treatment-intolerant or nonresponsive to standard therapy
Required Medical Information	Documentation of use in combination with bedaquiline and linezolid.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an infectious disease specialist.
Coverage Duration	Request will be authorized for 26 weeks.
Other Criteria	Documentation of prior trial of or medical reason for not using first-line TB regimen containing isoniazid and rifampin.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

PREVYMIS

Products Affected

- PREVYMIS ORAL

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be a hematologist, oncologist, infectious disease, or transplant specialist.
Coverage Duration	Request will be authorized for 6 months.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

PROMACTA

Products Affected

- *eltrombopag olamine oral packet 12.5 mg, 25 mg*
- *eltrombopag olamine oral tablet 12.5 mg, 25 mg, 50 mg, 75 mg*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	For chronic immune (idiopathic) thrombocytopenia (ITP): Documented baseline platelet count less than 30,000 cells/ microL. For severe aplastic anemia: Documentation of baseline platelet count less than 20,000 cells/microL OR platelet count less than 30,000 cells/microL with bleeding OR reticulocyte count less than 20,000 cells/microL OR absolute neutrophil count less than 500 cells/microL. For thrombocytopenia in patients with Hepatitis C infection: documented baseline platelet count less than 75,000 cells/microL.
Age Restrictions	N/A
Prescriber Restrictions	Specialist for submitted diagnosis.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	For chronic immune (idiopathic) thrombocytopenia (ITP): Trial of, contraindication to, or medical reason for not using glucocorticosteroids. For severe aplastic anemia: Trial of, contraindication to, or medical reason for not using at least one immunosuppressive agent.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

PYRUKYND

Products Affected

- PYRUKYND
- PYRUKYND TAPER PACK

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	For new starts: 1) documentation of diagnosis and 2) baseline hemoglobin level. For continuation of therapy or reauthorization: documentation of clinical improvement (e.g. reduction in number of blood transfusions, or increase or stabilization in hemoglobin level). If the criteria are not met, may authorize up to 14 days of a Pyrukynd Taper Pack to allow for tapering.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a hematologist.
Coverage Duration	New starts: 6 mo. Cont of therapy or reauth: end of contract yr. Denial: 14 days for dose tapering.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

RADICAVA

Products Affected

- RADICAVA ORS
- RADICAVA ORS STARTER KIT

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be a neurologist.
Coverage Duration	New starts: 6 months. Cont. of therapy or reauthorization: until end of contract year.
Other Criteria	For new starts: 1) documentation of ALS functional rating scale (ALSFRS-R) score and 2) documentation that the member has been on riluzole, is beginning therapy as an adjunct to treatment with Radicava, or provider has provided a medical reason why patient is unable to use riluzole. For continuation of therapy or reauthorization: documentation from provider of clinical stabilization in symptoms (e.g. stabilization of ALS functional rating scale (ALSFRS-R) score).
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

RAVICTI

Products Affected

- RAVICTI

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Provider is a geneticist, metabolic specialist, gastroenterologist, hepatologist, or liver transplant specialist.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	Trial of, contraindication to, or medical reason for not using sodium phenylbutyrate.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

RECORLEV

Products Affected

- RECORLEV

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Specialist for submitted diagnosis.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	Trial of, contraindication to, or medical reason for not using ketoconazole tablets.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

RELISTOR

Products Affected

- RELISTOR ORAL
- RELISTOR SUBCUTANEOUS SOLUTION
- RELISTOR SUBCUTANEOUS SOLUTION PREFILLED SYRINGE

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	Patient must have documented trial of or medical reason for not using the following: 1) lubiprostone, AND 2) lactulose AND 3) Movantik. Additionally, patient must have a medical reason for not being able to use oral Relistor in order to receive Relistor injection.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

REXULTI

Products Affected

- REXULTI

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	For schizophrenia: trial of, contraindication to, or medical reason for not using two generic antipsychotics. For major depressive disorder: trial of, contraindication to, or medical reason for not using to two generic antidepressants.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

REZUROCK

Products Affected

- REZUROCK

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be a hematologist, oncologist, or transplant specialist.
Coverage Duration	New starts: 3 months. Cont. of therapy or reauthorization: until end of contract year.
Other Criteria	For new starts: documented trial of, contraindication to, or medical reason for not using at least two lines of systemic immunosuppressive therapy (e.g. corticosteroids, tacrolimus, mycophenolate mofetil, Imbruvica, or Jakafi), one of which must be a systemic corticosteroid. For continuation of therapy or re-authorization: documentation of clinical benefit from use of the drug (i.e. symptom improvement, reduction in corticosteroid dose).
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

RINVOQ

Products Affected

- RINVOQ
- RINVOQ LQ

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Specialist for submitted diagnosis.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	For RA: Trial of, medical reason for not using, or contraindication to 1 disease modifying antirheumatic drug (DMARD) (methotrexate, leflunomide, or sulfasalazine) and 1 tumor necrosis factor (TNF) blocker (Enbrel, Hadlima, or Humira). For PsA: Trial of, medical reason for not using, or contraindication to 1 TNF blocker (Enbrel, Hadlima, or Humira). For atopic dermatitis: trial of, contraindication to, or medical reason for not using: 1) topical tacrolimus or pimecrolimus and 2) Eucrisa. For ankylosing spondylitis: Trial of, medical reason for not using, or contraindication to naproxen and 1 TNF blocker (Enbrel, Hadlima, or Humira). For UC: Trial of, medical reason for not using, or contraindication to 1 TNF blocker (ex. Enbrel, Hadlima, or Humira). For non radiographic axial spondyloarthritis: Trial of, medical reason for not using, or contraindication to naproxen. For Crohns Disease: trial of, medical reason for not using, or contraindication to 1 TNF blocker. For pJIA: Trial of, medical reason for not using, or contraindication to 1 TNF blocker (Enbrel, Hadlima, or Humira). For Giant cell arteritis: trial of, medical reason for not using, or contraindication to at least 1 corticosteroid (i.e. prednisone, methylprednisolone).
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

PA Criteria	Criteria Details
Part B Prerequisite	No

RUFINAMIDE

Products Affected

- *rufinamide oral suspension*
- *rufinamide oral tablet*

PA Criteria	Criteria Details
Exclusion Criteria	History of familial Short QT syndrome
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be a neurologist.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	Trial of, contraindication to, or medical reason for not using one alternative generic anticonvulsant for appropriate indications.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

RYKINDO

Products Affected

- RYKINDO

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	The member has a documented history of receiving oral risperidone without any clinically significant side effects.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	Trial of, contraindication to, or medical reason (e.g. intolerance, hypersensitivity or contraindication) for not using at least two of the following: Abilify Maintena, Abilify Asimtufii, Invega Sustenna, Invega Trinza, or Risperidone Microspheres ER.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

RYLAZE

Products Affected

- RYLAZE

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be an oncologist, hematologist, or specialist for submitted diagnosis.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

SAPROPTERIN

Products Affected

- *sapropterin dihydrochloride oral packet*
- *sapropterin dihydrochloride oral tablet*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	For new starts: documentation of elevated baseline phenylalanine levels. Continuation of therapy or reauthorization: prescriber attests the member has improvement in phenylalanine levels from baseline.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	New starts will be authorized for 3 months. Cont of therapy or reauth until end of contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

SECUADO

Products Affected

- SECUADO

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	Trial of, contraindication to, or medical reason for not using to one generic antipsychotics.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

SEROSTIM

Products Affected

- SEROSTIM SUBCUTANEOUS
SOLUTION RECONSTITUTED 4 MG, 5
MG, 6 MG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a HIV specialist, gastroenterologist, nutritional support specialist or ID specialist.
Coverage Duration	Request will be authorized for 12 weeks.
Other Criteria	For initial starts for HIV wasting/cachexia: 1) Member must be on anti-retroviral therapy and 2) Trial of, contraindication to or medical reason for not using megestrol or dronabinol and 3) Alternative causes of wasting have been ruled out (diarrhea, malignancies, inadequate caloric intake, etc)
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

SIGNIFOR

Products Affected

- SIGNIFOR

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Member is not a candidate for surgery or surgery was not curative.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an endocrinologist.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

SILDENAFIL ORAL

Products Affected

- *sildenafil citrate oral suspension reconstituted*
- *sildenafil citrate oral tablet 20 mg*

PA Criteria	Criteria Details
Exclusion Criteria	Documentation of concurrent nitrate or Adempas use.
Required Medical Information	Pulmonary Arterial Hypertension: Documentation of pulmonary arterial hypertension (PAH) WHO Group I and PAH Functional Class. Reviewer will verify available patient claim history to confirm patient is not using nitrates or Adempas. Secondary Raynaud's phenomenon- Initial: [Note: documentation required] (1) diagnosis of secondary Raynaud's phenomenon AND (2) diagnosis of primary condition which Raynaud's phenomenon is secondary to (e.g., lupus, scleroderma, rheumatoid arthritis, Sjogren's syndrome, thyroid disease). Continuation of therapy: patient has positive clinical response to treatment.
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be a pulmonologist, cardiologist, or rheumatologist.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	For sildenafil suspension: Documentation of trial of, contraindication to, or medical reason for not using sildenafil tablet.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

SILIQ

Products Affected

- SILIQ

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Specialist for submitted diagnosis.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	For psoriasis: Either 1) Trial of, medical reason for not using, or contraindication to 2 of the following therapies: Stelara, Skyrizi, Tremfya, Enbrel, Hadlima, or Humira, or 2) If utilized within the past 120 days, approve for continuation of therapy.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

SIMPONI

Products Affected

- SIMPONI SUBCUTANEOUS SOLUTION AUTO-INJECTOR
- SIMPONI SUBCUTANEOUS SOLUTION PREFILLED SYRINGE

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Specialist for submitted diagnosis.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	For ankylosing spondylitis: Either 1) Trial of, medical reason for not using, or contraindication to 2 of the following therapies: Enbrel, Humira, Hadlima, Rinvoq or Xeljanz, or 2) If utilized within the past 120 days, approve for continuation of therapy. For PsA: Either 1) Trial of, medical reason for not using, or contraindication to 2 of the following therapies: Stelara, Skyrizi, Tremfya, Xeljanz, Rinvoq, Enbrel, Hadlima, or Humira, or 2) If utilized within the past 120 days, approve for continuation of therapy. For RA: Either 1) Trial of, medical reason for not using, or contraindication to 2 of the following: Enbrel, Humira, Hadlima, Rinvoq or Xeljanz or 2) If utilized within the past 120 days, approve for continuation of therapy. For UC: Either 1) Trial of, medical reason for not using, or contraindication to 2 of the following: Skyrizi, Humira, Hadlima, Rinvoq, Stelara, Tremfya, or Xeljanz or 2) If utilized within the past 120 days, approve for continuation of therapy.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

SIRTURO

Products Affected

- SIRTURO

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Documentation (consistent with pharmacy claims data, OR for new members to the health plan consistent with medical chart history) that the member is currently taking 3 additional antimycobacterial drugs in combination to treat MDR-TB.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an infectious disease specialist.
Coverage Duration	Request will be authorized for 24 weeks.
Other Criteria	Documentation of prior trial of or medical reason for not using first-line TB regimen containing isoniazid and rifampin.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

SKYRIZI

Products Affected

- SKYRIZI
- SKYRIZI PEN

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Specialist for submitted diagnosis.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	For PsA or psoriasis: approve. For Crohns Disease: Either 1) Trial of, medical reason for not using (i.e. severe Crohns disease), or contraindication to 1 of the following: mercaptopurine, azathioprine, sulfasalazine, methotrexate or corticosteroid (e.g., prednisone, methylprednisolone) or 2) If utilized within the past 120 days, approve for continuation of therapy. For UC: Trial of, medical reason for not using, or contraindication to 1 of the following conventional therapies: mercaptopurine, an aminosalicylate (i.e. mesalamine, sulfasalazine, azathioprine), or a corticosteroid (i.e. prednisone, methylprednisolone).
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

SODIUM OXYBATE

Products Affected

- *sodium oxybate*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be a sleep specialist, pulmonologist, or neurologist.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	For somnolence associated with narcolepsy: trial of, contraindication to, or medical reason for not using a CNS stimulant (e.g. methylphenidate, modafinil, armodafinil, etc.). For cataplexy associated with narcolepsy, approve.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

SODIUM PHENYLBUTYRATE

Products Affected

- *sodium phenylbutyrate oral powder 3 gm/tsp*
- *sodium phenylbutyrate oral tablet*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Provider is a geneticist, metabolic specialist, gastroenterologist, hepatologist, or liver transplant specialist.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

SOFOSBUVIR/VELPATASVIR

Products Affected

- SOFOSBUVIR-VELPATASVIR

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Detectable HCV RNA viral load prior to treatment within 6 months of request. In addition, documentation of treatment history, and if cirrhotic, documentation of compensated or decompensated cirrhosis.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized for 12-24 weeks based on AASLD-IDSA guidelines
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

SOMAVERT

Products Affected

- SOMAVERT

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	For new starts for acromegaly: pt meets one of the following (1) inadequate response to surgery and/or radiotherapy OR (2) pt is not an appropriate candidate for surgery and/or radiotherapy OR (3) pt is experiencing negative effects due to tumor size (ex: optic nerve compression). Continuation of therapy or reauthorization: documentation of clinical improvement with therapy.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an endocrinologist.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

SOTYKTU

Products Affected

- SOTYKTU

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Specialist for submitted diagnosis.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	For moderate to severe psoriasis: Either 1) Trial of, medical reason for not using, or contraindication to 2 of the following therapies: Stelara, Skyrizi, Tremfya, Enbrel, Hadlima, or Humira, or 2) If utilized within the past 120 days, approve for continuation of therapy.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

STELARA

Products Affected

- STELARA INTRAVENOUS
- STELARA SUBCUTANEOUS SOLUTION 45 MG/0.5ML
- STELARA SUBCUTANEOUS SOLUTION PREFILLED SYRINGE

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Specialist for submitted diagnosis.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	For Crohns Disease: Either 1) Trial of, medical reason for not using (i.e. severe Crohns disease), or contraindication to 1 of the following: mercaptopurine, azathioprine, methotrexate, sulfasalazine, or corticosteroid (e.g., prednisone, methylprednisolone) or 2) If utilized within the past 120 days, approve for continuation of therapy. For psoriasis: Approve. For PsA: Approve. For UC: Either 1) Trial of, medical reason for not using, or contraindication to 1 of the following conventional therapies: mercaptopurine, an aminosalicylate (i.e. mesalamine, sulfasalazine, azathioprine), or a corticosteroid (i.e. prednisone, methylprednisolone) or 2) If utilized within the past 120 days, approve for continuation of therapy.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

SUCRAID

Products Affected

- SUCRAID

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	For new starts: documentation of diagnosis of congenital sucrase-isomaltase deficiency. For continuation of therapy or reauthorization: Prescriber attests that member has obtained a clinical benefit (e.g. fewer total stools, greater number of hard and formed stools, fewer watery and soft stools, decrease in breath hydrogen output)
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	New starts will be authorized for 3 months. Cont of therapy or reauth until end of contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

SYMDEKO

Products Affected

- SYMDEKO

PA Criteria	Criteria Details
Exclusion Criteria	Combination use with Kalydeco, Orkambi, or Trikafta.
Required Medical Information	Documentation of CFTR gene that is responsive to tezacaftor-ivacaftor treatment.
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be a pulmonologist or an expert in the treatment of cystic fibrosis.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

SYMLIN

Products Affected

- SYMLINPEN 120 SUBCUTANEOUS SOLUTION PEN-INJECTOR
- SYMLINPEN 60 SUBCUTANEOUS SOLUTION PEN-INJECTOR

PA Criteria	Criteria Details
Exclusion Criteria	Patient has confirmed gastroparesis.
Required Medical Information	For new starts: HbA1C values within 90 days of request is greater than or equal to 7% despite receiving insulin therapy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	Trial of, contraindication to, or medical reason for not using two alternative anti-diabetic agents.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

SYNAREL

Products Affected

- SYNAREL

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	Trial of, contraindication to, or medical reason for not using the following concurrently for endometriosis: analgesic pain reliever (e.g. NSAIDs, COX-2 inhibitors) AND either combined estrogen-progestin oral contraceptive, progestin (e.g. medroxyprogesterone acetate, norethindrone), OR gonadotropin-releasing hormone (GnRH) agonists (e.g. Lupron Depot)
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

TADALAFIL

Products Affected

- *tadalafil (pah)*
- TADLIQ

PA Criteria	Criteria Details
Exclusion Criteria	Documentation of concurrent nitrate or Adempas use.
Required Medical Information	Documentation of pulmonary arterial hypertension (PAH) WHO Group I and PAH Functional Class. Reviewer will verify available patient claim history to confirm patient is not using nitrates or Adempas.
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be a pulmonologist or cardiologist.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	For Tadliq: Documentation of trial of, contraindication to, or medical reason for not using tadalafil tablets.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

TADALAFIL, BPH

Products Affected

- *tadalafil oral tablet 5 mg*

PA Criteria	Criteria Details
Exclusion Criteria	Diagnosis of erectile dysfunction
Required Medical Information	Diagnosis of Benign prostatic hyperplasia (BPH) required AND trial of, contraindication to, or medical reason for not using an alpha blocker (e.g. tamsulosin, terazosin).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

TALTZ

Products Affected

- TALTZ

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Specialist for submitted diagnosis.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	For ankylosing spondylitis: Either 1) Trial of, medical reason for not using, or contraindication to 2 of the following therapies: Enbrel, Humira, Hadlima, Rinvoq or Xeljanz, or 2) If utilized within the past 120 days, approve for continuation of therapy. For non-radiographic axial spondyloarthritis: approve. For psoriasis: Either 1) Trial of, medical reason for not using, or contraindication (e.g., safety concerns, not indicated for patient's age) to 2 of the following therapies: Stelara, Skyrizi, Tremfya, Enbrel, Hadlima, or Humira, or 2) If utilized within the past 120 days, approve for continuation of therapy. For PsA: Either 1) Trial of, medical reason for not using, or contraindication to 2 of the following therapies: Stelara, Skyrizi, Tremfya, Xeljanz, Rinvoq, Enbrel, Hadlima, or Humira, or 2) If utilized within the past 120 days, approve for continuation of therapy.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

TARPEYO

Products Affected

- TARPEYO

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	For new starts: attestation that member has 1) Diagnosis of primary immunoglobulin A nephropathy (IgAN) and 2) at risk of disease progression. Member has an estimated glomerular filtration rate (eGFR) greater than or equal to 35 mL/min/1.73 m(2) and proteinuria. For continuation of therapy: documentation that member has been on Tarpeyo for less than 9 months. For reauthorizations: Requests will not be allowed as the safety and efficacy of subsequent courses of Tarpeyo have not been established.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a nephrologist.
Coverage Duration	Request will be authorized for 9 months.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

TAVNEOS

Products Affected

- TAVNEOS

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a rheumatologist or hematologist.
Coverage Duration	New starts will be authorized for 6 months. Cont of therapy or reauth until end of contract year.
Other Criteria	For new starts: 1) Prescriber attests that Tavneos will be prescribed in combination with corticosteroids AND cyclophosphamide unless there is documented trial of, contraindication to, or medical reason for not using these therapies. 2) Documentation of baseline Birmingham Vasculitis Activity Score (BVAS) score 3) Prescriber attestation that the patient will have liver function tests before treatment (ALT, AST, alkaline phosphate, and total bilirubin) and every 4 weeks after start of therapy for the first 6 months of treatment 4) Prescriber attestation that the patient has been screened for and does not have active hepatitis B virus (HBV) infection at baseline. For continuation of therapy or reauthorization: 1) Documentation of remission (BVAS score of 0) OR improvement in BVAS score 2) Prescriber attestation that patient has no abnormality in liver function tests (abnormality: ALT or AST greater than 3 times the upper limit of normal and bilirubin greater than 2 times the upper limit of normal) 3) Prescriber attestation that patient has no active HBV infection.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

TECENTRIQ HYBREZA

Products Affected

- TECENTRIQ HYBREZA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be an oncologist or specialist for submitted diagnosis.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

TEFLARO

Products Affected

- TEFLARO

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Documentation of a consultation with an infectious disease specialist.
Coverage Duration	Request will be authorized for 14 days.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

TEPEZZA

Products Affected

- TEPEZZA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Documentation of moderate to severe active thyroid eye disease (TED) as evidenced by one or more of the following: a) eyelid retraction greater than or equal to 2 mm, b) moderate or severe soft tissue involvement, c) exophthalmos greater than or equal to 3 mm above normal for race and sex or d) periodic or constant diplopia OR Documentation of chronic TED with one or more of the following: a) a 3 mm or greater increase in exophthalmos from before diagnosis of TED, b) exophthalmos greater than or equal to 3 mm above normal for race and sex.
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be an ophthalmologist or endocrinologist
Coverage Duration	Request will be authorized for 6 months.
Other Criteria	For active TED: member has had a trial and failure, contraindication to, or medical reason for not using oral or IV glucocorticoids.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

TERIPARATIDE

Products Affected

- BONSITY
- TERIPARATIDE SUBCUTANEOUS SOLUTION PEN-INJECTOR 560 MCG/2.24ML

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Documentation showing patient falls into one of the following categories: a bone mineral density (BMD) value consistent with osteoporosis (i.e., T-scores equal to or less than -2.5) or patient has had an osteoporotic fracture or patient has T-scores from -1.5 to -2.5 at the femoral neck or spine, and a 10-year probability of hip fracture greater than or equal to 3% or a 10-year probability of any major osteoporosis-related fracture greater than or equal to 20% based on the United States-adapted FRAX model.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	In addition, the following criteria is also applicable: Trial of, medical reason for not using, or contraindication to an oral bisphosphonate and Prolia
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

THIOLA

Products Affected

- *tiopronin oral*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

TIGECYCLINE

Products Affected

- *tigecycline*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	(1) Patient must have documented diagnosis of one of the following infections: (a) complicated skin and skin structure infection, (b) complicated intraabdominal infection, (c) community-acquired pneumonia AND (2) Trial of, contraindication to, or medical reason (e.g. intolerance, hypersensitivity or contraindication) for not using preferred first-line antibiotics.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an infectious disease specialist.
Coverage Duration	Request will be authorized for 14 days.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

TOLVAPTAN

Products Affected

- *tolvaptan*

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with strong CYP3A4 inhibitors (i.e. clarithromycin, ketoconazole, itraconazole, ritonavir, lopinavir-ritonavir, indinavir-ritonavir, indinavir, nelfinavir, saquinavir, nefazodone, conivaptan, and telithromycin).
Required Medical Information	Reviewer will verify available patient claim history to confirm patient is not using a strong CYP3A4 inhibitor (i.e. clarithromycin, ketoconazole, itraconazole, ritonavir, lopinavir-ritonavir, indinavir-ritonavir, indinavir, nelfinavir, saquinavir, nefazodone, conivaptan, and telithromycin).
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a cardiologist, endocrinologist, hepatologist, or nephrologist.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

TOPICAL ANTINEOPLASTIC RETINOIDS

Products Affected

- *bexarotene*
- PANRETIN

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

TOPICAL TESTOSTERONE

Products Affected

- *testosterone transdermal gel 1.62 %, 12.5 mg/act (1%), 20.25 mg/1.25gm (1.62%), 20.25 mg/act (1.62%), 25 mg/2.5gm (1%), 40.5 mg/2.5gm (1.62%), 50 mg/5gm (1%)*
- *testosterone transdermal solution*

PA Criteria	Criteria Details
Exclusion Criteria	Patient has history of prostate cancer or breast cancer.
Required Medical Information	New starts of topical testosterone therapy for hypogonadism must have both of the following characteristics of hypogonadism: 1) symptoms associated with hypogonadism (e.g. unexplained mild anemia, low libido, decreased energy, etc.) 2) Two separate instances of low serum total or free testosterone taken in the morning, as defined by the lab reference range.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

TRANSDERMAL LIDOCAINE

Products Affected

- *lidocaine external patch 5 %*
- ZTLIDO

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	If the request is for the product ZTlido, must provide medical reason for not being able to use generic lidocaine 5% patch
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

TREMFYA

Products Affected

- TREMFYA CROHNS INDUCTION
- TREMFYA ONE-PRESS
SUBCUTANEOUS SOLUTION PEN-
INJECTOR
- TREMFYA PEN
- TREMFYA SUBCUTANEOUS
SOLUTION PREFILLED SYRINGE
- TREMFYA-CD/UC INDUCTION

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Specialist for submitted diagnosis.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	For UC: Approve. Crohn's Disease (CD): Trial of, medical reason for not using (i.e. severe Crohn's disease), or contraindication to 1 of the following: mercaptopurine, azathioprine, sulfasalazine, methotrexate or corticosteroid (e.g., prednisone, methylprednisolone).
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

TRIENTINE

Products Affected

- CUVRIOR
- *trientine hcl oral capsule 250 mg*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	If the request is for Cuvrior for new starts, member must have trial of, contraindication to, or medical reason for not using trientine hydrochloride.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

TRIKAFTA

Products Affected

- TRIKAFTA

PA Criteria	Criteria Details
Exclusion Criteria	Combination use with Kalydeco, Orkambi, or Symdeko.
Required Medical Information	Documentation of CFTR gene that is responsive to elexacaftor-tezacaftor-ivacaftor treatment.
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be a pulmonologist or an expert in the treatment of cystic fibrosis.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

TYMLOS

Products Affected

- TYMLOS

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Documentation showing patient falls into one of the following categories: a bone mineral density (BMD) value consistent with osteoporosis (i.e., T-scores equal to or less than -2.5) or patient has had an osteoporotic fracture or patient has T-scores from -1.5 to -2.5 at the femoral neck or spine, and a 10-year probability of hip fracture greater than or equal to 3% or a 10-year probability of any major osteoporosis-related fracture greater than or equal to 20% based on the United States-adapted FRAX model.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	The following criteria is also applicable: 1) trial of, contraindication to, or medical reason for not using an oral bisphosphonate and Prolia, and 2) therapy does not exceed 2 years.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

TYVASO

Products Affected

- TYVASO DPI MAINTENANCE KIT
INHALATION POWDER 16 MCG, 32 MCG, 48 MCG, 64 MCG
- TYVASO DPI TITRATION KIT
INHALATION POWDER 16 & 32 & 48 MCG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be a pulmonologist or cardiologist.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	For the treatment of pulmonary arterial hypertension (PAH): 1) documentation of PAH WHO Group I classification and PAH Functional Class and 2) trial of, contraindication to, or medical reason for not using a generic phosphodiesterase inhibitor and a generic endothelin receptor antagonist. For the treatment of pulmonary hypertension associated with interstitial lung disease (PH-ILD, WHO Group 3): documentation of PH-ILD and PAH Functional Class.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

UPTRAVI

Products Affected

- UPTRAVI ORAL
- UPTRAVI TITRATION

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Documentation of pulmonary arterial hypertension (PAH) WHO Group I and PAH Functional Class.
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be a pulmonologist or cardiologist.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	Trial of, contraindication to, or medical reason for not using a generic phosphodiesterase inhibitor and a generic endothelin receptor antagonist.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

UZEDY

Products Affected

- UZEDY

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	The member has a documented history of receiving oral risperidone without any clinically significant side effects.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	Trial of, contraindication to, or medical reason (e.g. intolerance, hypersensitivity or contraindication) for not using at least two of the following: Abilify Maintena, Abilify Asimtufii, Invega Sustenna, Invega Trinza, or Risperdion Microsphere ER.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

VALCHLOR

Products Affected

- VALCHLOR

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or dermatologist.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	Trial of, contraindication to, or medical reason for not being able to use one of the following: a topical corticosteroids or a topical retinoids.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

VEMLIDY

Products Affected

- VEMLIDY

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	For new starts: attestation that member has been tested for HIV infection. If member is HIV-positive, Vemlidy is not used alone.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

VENTAVIS

Products Affected

- VENTAVIS

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Documentation of pulmonary arterial hypertension (PAH) WHO Group I classification and PAH Functional Class.
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be a pulmonologist or cardiologist.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

VEOZAH

Products Affected

- VEOZAH

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Initial: (1) Documented diagnosis of moderate to severe vasomotor symptoms due to menopause AND (2) Trial of, contraindication to, or medical reason (e.g. intolerance, hypersensitivity or contraindication) for not using a hormonal therapy (e.g., estradiol, oral Premarin, Prempro). Reauthorization: (1) Documentation of positive clinical response to therapy (e.g., decrease in frequency or severity of vasomotor symptoms from baseline)
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

VIGABATRIN

Products Affected

- *vigabatrin*
- VIGAFYDE

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	For infantile spasms or West syndrome, the request will be approved. For diagnosis of refractory complex partial seizures: 1) documentation of diagnosis, and 2) attestation the member is currently receiving another antiepileptic drug, and 3) attestation the member has experienced treatment failure from two alternative antiepileptic agents.
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be a neurologist.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

VIJOICE

Products Affected

- VIJOICE

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	For new starts, all of the following must be included: 1) Documentation of genetic testing confirming diagnosis AND 2) Member has at least one target lesion identified on imaging AND 3) Prescriber attests the patient's condition is severe or life-threatening and necessitates systemic treatment. For continuation of therapy or reauthorization, attestation of a positive clinical response (i.e. reduction in the sum of measurable target lesion volume, absence of progression of non-target lesions, absence of any new lesions, etc.).
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a geneticist, dermatologist, vascular surgeon, hematologist/oncologist, or other specialist in the treatment of PIK3CA-Related Overgrowth Spectrum(PROs).
Coverage Duration	New starts will be authorized for 6 months. Cont of therapy or reauth until end of contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

VMAT-2 INHIBITORS

Products Affected

- AUSTEDO
- AUSTEDO PATIENT TITRATION KIT
- AUSTEDO XR
- AUSTEDO XR PATIENT TITRATION ORAL TABLET EXTENDED RELEASE THERAPY PACK 12 & 18 & 24 & 30 MG
- INGREZZA ORAL CAPSULE
- INGREZZA ORAL CAPSULE SPRINKLE
- INGREZZA ORAL CAPSULE THERAPY PACK
- *tetrabenazine*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be a neurologist, clinical geneticist, or psychiatrist.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	If the request is for tetrabenazine, Ingrezza or Ingrezza Sprinkle, request will be approved. If the request is for Austedo or Austedo XR, the member must have trial of or medical reason for not using tetrabenazine. Reauthorization: Confirmation of improvement in tardive dyskinesia symptoms or chorea associated with Huntington disease symptoms.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

VORICONAZOLE

Products Affected

- *voriconazole intravenous*

PA Criteria	Criteria Details
Exclusion Criteria	Non-Part D indications.
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized for 6 months.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

VOSEVI

Products Affected

- VOSEVI

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Detectable HCV RNA viral load prior to treatment within 6 months of request. In addition, documentation of treatment history, and if cirrhotic, documentation of compensated or decompensated cirrhosis.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized for 12 weeks as per AASLD-IDSA guidance.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

VOWST

Products Affected

- VOWST

PA Criteria	Criteria Details
Exclusion Criteria	Treatment of Clostridioides difficile infection (CDI)
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	If all the criteria are met, the request will be approved for 1 month
Other Criteria	Diagnosis of at least 1 recurrent episode of CDI
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

WEGOVY

Products Affected

- WEGOVY SUBCUTANEOUS SOLUTION AUTO-INJECTOR 0.25 MG/0.5ML, 1.7 MG/0.75ML, 2.4 MG/0.75ML, 0.5 MG/0.5ML, 1

PA Criteria	Criteria Details
Exclusion Criteria	The member has an indication of only weight reduction or maintenance for overweight or obesity. The member has concurrent use of any GLP-1 receptor agonist. The member has a personal history of medullary thyroid carcinoma. The member has Multiple Endocrine Neoplasia syndrome type 2. For a diagnosis of reduction of the risk of major cardiovascular events in adults with established cardiovascular disease and either obesity or overweight: the member has a personal history of Type 1 or Type 2 diabetes.
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	Reduction of the risk of major adverse cardiovascular events in adults with established cardiovascular disease and either obesity or overweight - For new starts: The member has an indication for reducing the risk of adverse cardiovascular events (cardiovascular death, non-fatal myocardial infarction, or non-fatal stroke) in adults with established cardiovascular disease. Documentation demonstrates patient has established cardiovascular disease (i.e., prior myocardial infarction, prior stroke, symptomatic peripheral arterial disease). Documentation is provided that the patient is overweight or obese (defined as a BMI of greater than or equal to 27 kg/m ²). Documentation is provided that the patients Hb A1c is less than or equal to 6.5%. For continuation of therapy or reauthorization: Documentation is provided that the patients Hb A1c is less than or equal to 6.5%. Patient continues to not have Type 1 or Type 2 diabetes. Treatment of noncirrhotic metabolic dysfunction-associated steatohepatitis (MASH) - For new starts: The member has a documented diagnosis of MASH with

PA Criteria	Criteria Details
	documentation of moderate to advanced liver fibrosis (fibrosis stage 2 or 3) as evidenced by liver biopsy, biomarker testing (AST to platelet ratio index score, fibrosis 4 score, NAFLD fibrosis score, etc.), transient elastography or ultrasonography. For continuation of therapy or reauthorization: Documentation is provided that the patient continues to have a fibrosis stage of less than or equal to 3.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

WHITE BLOOD CELL STIMULATORS

Products Affected

- FULPHILA
- FYLNETRA
- LEUKINE INJECTION SOLUTION RECONSTITUTED
- NEULASTA ONPRO
- NEULASTA SUBCUTANEOUS SOLUTION PREFILLED SYRINGE
- NIVESTYM
- NYVEPRIA
- RELEUKO SUBCUTANEOUS
- STIMUFEND
- UDENYCA
- UDENYCA ONBODY
- ZARXIO
- ZIEXTENZO

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	For new starts for Neulasta, Udenyca, Ziextenzo, Stimufend and Nyvepria: documentation of trial of, contraindication to, or medical reason for not using Fylnetra and Fulphila. Continuation of therapy or re-authorization criteria: diagnosis of chronic neutropenia or a medical reason for continued need for GCSF.
Age Restrictions	N/A
Prescriber Restrictions	Specialist for submitted diagnosis.
Coverage Duration	New starts will be authorized for 4 months. Cont of therapy or reauth until end of contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

XATMEP

Products Affected

- XATMEP

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be an oncologist or rheumatologist.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

XDEM VY

Products Affected

- XDEM VY

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

XELJANZ

Products Affected

- XELJANZ
- XELJANZ XR

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Specialist for submitted diagnosis.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	For ankylosing spondylitis: Trial of, medical reason for not using, or contraindication to naproxen and 1 TNF blocker (Enbrel, Hadlima, or Humira) For pJIA: Trial of, medical reason for not using, or contraindication to 1 of the following DMARDs: methotrexate or leflunomide and 1 TNF blocker (Enbrel, Hadlima, or Humira). For PsA: Trial of, medical reason for not using, or contraindication to 1 TNF blocker (Enbrel, Hadlima, or Humira). For RA: Trial of, medical reason for not using, or contraindication to 1 disease modifying antirheumatic drug (DMARD) (methotrexate, leflunomide, or sulfasalazine) and 1 tumor necrosis factor (TNF) blocker (Enbrel, Hadlima, or Humira). For UC: Trial of, medical reason for not using, or contraindication to 1 of the following conventional therapies: mercaptopurine, an aminosalicylate (i.e. mesalamine, sulfasalazine, azathioprine), or a corticosteroid (i.e. prednisone, methylprednisolone) and Humira or Hadlima.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

XERMELO

Products Affected

- XERMELO

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a gastroenterologist or an oncologist.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	For new starts: 1) Attestation that diarrhea is inadequately controlled by stable dose of SSA therapy for at least three months. For continuation of therapy or reauthorization: 1) documentation of positive clinical response to xermelo and 2) Attestation to continue to be used in combination with SSA.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

XGEVA

Products Affected

- OSENVELT
- WYOST
- XGEVA

PA Criteria	Criteria Details
Exclusion Criteria	Patients with baseline hypocalcemia
Required Medical Information	New starts: Serum calcium levels. Reauthorization criteria for malignant hypercalcemia: albumin-adjusted serum calcium level below 12.5mg/dl within 30 days of request.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

XIFAXAN

Products Affected

- XIFAXAN

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	For HE: gastroenterologist or hepatologist. For IBS-D: gastroenterologist.
Coverage Duration	For HE: contract year. For IBSD: 14 days (cannot exceed 3 courses of 14 days each). For TD: 3 days.
Other Criteria	For diagnosis of hepatic encephalopathy (HE): trial of, contraindication to, or medical reason for not using lactulose. For diagnosis of irritable bowel syndrome with diarrhea (IBSD): No more than 3 courses of 14 days each. For travelers diarrhea (TD) caused by noninvasive strains of E. Coli (with no bloody stools or fever): patient must be intolerant to or must have had a trial of at least 3 days of one of the following agents: ciprofloxacin, ofloxacin, levofloxacin or azithromycin.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

XOLAIR

Products Affected

- XOLAIR

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be a pulmonologist, allergist, immunologist, dermatologist, or otolaryngologist.
Coverage Duration	New starts will be authorized for 6 months. Cont of therapy or reauth until end of contract year.
Other Criteria	New starts for moderate to severe persistent allergic asthma: 1) Evidence of specific allergic sensitivity confirmed by positive skin test (i.e. prick/puncture test) or blood test (i.e. radioallergosorbent test) for a specific IgE or in vitro reactivity to a perennial aeroallergen, AND 2) Pretreatment serum IgE levels greater than 30 IU/mL, AND 3) Symptoms are not adequately controlled with high-dose inhaled corticosteroid (ICS) plus additional controller medication (ie. long-acting B2 agonist) for at least 3 months, or there is a medical reason for not using these drugs. Continuation of therapy or reauthorization criteria for moderate to severe persistent allergic asthma: 1) Reduction in asthma exacerbation resulting in systemic steroid use and/or hospitalization, OR 2) Reduction of rescue inhaler use, OR 3) Documentation of improvement in pulmonary function tests since baseline (prior to initiation of Xolair). New starts for chronic idiopathic urticaria: 1) inadequate symptomatic relief despite trial of two weeks of two different oral antihistamine therapies (unless contraindicated), AND 2) disease must be severe enough to warrant short term systemic corticosteroid therapy for management of urticaria. Continuation of therapy or reauthorization criteria for chronic idiopathic urticaria: 1) improvement from baseline of symptoms associated with urticaria within 6 months of Xolair use. New starts for nasal polyps: 1) currently using an intranasal corticosteroid, will be prescribed an intranasal corticosteroid with request, or has a medical reason for not using an intranasal corticosteroid.

PA Criteria	Criteria Details
	Continuation of therapy or reauthorization criteria for nasal polyps: 1) Documentation has been provided that demonstrates a clinical benefit (e.g. improvements in symptom severity, nasal polyp score [NPS], sino-nasal outcome test-22 [SNOT-22], nasal congestion score [NCS]) AND 2) continued use of intranasal corticosteroid, or has a medical reason for not using one. New starts for food allergy: 1) diagnosis of IgE-mediated food allergy 2) Xolair will be used in conjunction with food allergen avoidance. Continuation of therapy or reauthorization criteria for food allergy: 1) Documentation of clinical benefit
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

XOLREMDI

Products Affected

- XOLREMDI

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be an immunologist or a hematologist
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	For new starts: 1) A documented diagnosis of WHIM (warts, hypogammaglobulinemia, infections and myelokathexis) syndrome confirmed by genotype variant of chemokine receptor 4 (CXCR4) and absolute neutrophil count (ANC) of less than or equal to 400 cells/microliter or white blood cells (WBC) less than or equal to 400 cells/microliter and 2) Documentation of baseline ANC and absolute lymphocyte count (ALC). For renewal 1) Documentation or provider attestation of positive clinical response (i.e. improvement from baseline in ANC, WBC and/or ALC or reduced frequency, duration, or severity of infections, fewer warts, or improved or stabilized clinical signs and/or symptoms of WHIM).
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

XYWAV

Products Affected

- XYWAV

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be a sleep specialist, pulmonologist or a neurologist.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	For treatment of somnolence associated with narcolepsy, patient must have documentation of either trial of or a medical reason for being unable to use a CNS stimulant (e.g. methylphenidate, modafinil, armodafinil, etc.). For the treatment of cataplexy associated with narcolepsy or idiopathic hypersomnia, approve.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

YORVIPATH

Products Affected

- YORVIPATH

PA Criteria	Criteria Details
Exclusion Criteria	Diagnosis of acute post-surgical hypoparathyroidism.
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an endocrinologist.
Coverage Duration	New starts will be authorized for 6 months. Cont of therapy or reauth until end of contract year.
Other Criteria	For new starts: 1) Documented diagnosis of chronic hypoparathyroidism AND 2) Provider attestation that patient is currently receiving or has medical reason for not receiving calcium supplementation and active vitamin D treatment AND 3) An albumin-corrected serum calcium level of 7.8 mg/dL or greater. For reauthorization: Documentation of improvement in albumin-corrected serum calcium from baseline.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

YUTREPIA

Products Affected

- YUTREPIA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a cardiologist or a pulmonologist.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	For the treatment of pulmonary arterial hypertension (PAH): 1) documentation of PAH WHO Group I classification and PAH Functional Class and 2) trial of, contraindication to, or medical reason for not using a generic phosphodiesterase inhibitor and a generic endothelin receptor antagonist. For the treatment of pulmonary hypertension associated with interstitial lung disease (PH-ILD, WHO Group 3): documentation of PHILD and PAH Functional Class.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

ZEPBOUND

Products Affected

- ZEPBOUND SUBCUTANEOUS SOLUTION AUTO-INJECTOR

PA Criteria	Criteria Details
Exclusion Criteria	The member has an indication of only weight reduction or maintenance for overweight or obesity. The member has concurrent use of any GLP-1 receptor agonist. The member has a personal history of medullary thyroid carcinoma. The member has Multiple Endocrine Neoplasia syndrome type 2.
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a sleep disorder specialist, pulmonologist, ENT, or other provider specializing in obstructive sleep apnea.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	For new starts: The member has an indication for moderate to severe obstructive sleep apnea (OSA) in adults with obesity. Documentation of diagnosis of OSA through polysomnography (sleep study) with an apnea-hypopnea index of 15 or more events per hour, or five or more events per hour in the presence of symptoms (e.g., cognitive impairment, fatigue, insomnia, loud snoring) or cardiovascular comorbidities (e.g., hypertension, ischemic heart disease, previous stroke). Documentation is provided that the patient is obese (defined as a BMI of greater than or equal to 30 kg/m ²). For continuation of therapy: Documentation of positive response to treatment. Documentation member has achieved and/or maintained a decrease in weight since baseline.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID 25406

Last Update: 10/17/2025

ZEPOSIA

Products Affected

- ZEPOSIA
- ZEPOSIA 7-DAY STARTER PACK
- ZEPOSIA STARTER KIT ORAL CAPSULE THERAPY PACK 0.23MG & 0.46MG 0.92MG(21)

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Documentation of liver function tests (for new starts and for continuation of therapy or reauthorization)
Age Restrictions	N/A
Prescriber Restrictions	Specialist for submitted diagnosis.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	For multiple sclerosis: Trial of, contraindication to, or medical reason for not using two of the following: dalfampridine ER, dimethyl fumarate, fingolimod, glatiramer, glatopa, or teriflunomide. For ulcerative colitis: Either 1) Trial of, medical reason for not using, or contraindication Humira or 2) If utilized within the past 120 days, approve for continuation of therapy.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

ZILBRYSQ

Products Affected

- ZILBRYSQ

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be a neurologist, rheumatologist, or other appropriate specialist
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	Patient has tried and failed, a medical reason for not using, or has a contraindication to two (2) or more conventional therapies (i.e. pyridostigmine, corticosteroids, or non-steroidal immunosuppressive therapies)
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

ZTALMY

Products Affected

- ZTALMY

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a neurologist.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

ZURZUVAE

Products Affected

- ZURZUVAE

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	The member has a documented diagnosis of postpartum depression
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a psychiatrist or obstetrician/gynecologist
Coverage Duration	Request will be authorized until the end of the contract year
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

ZYPREXA RELPREVV

Products Affected

- ZYPREXA RELPREVV INTRAMUSCULAR SUSPENSION RECONSTITUTED 210 MG, 300 MG, 405 MG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	The member has a documented history of receiving oral olanzapine without any clinically significant side effects.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	Trial of, contraindication to, or medical reason (e.g. intolerance, hypersensitivity or contraindication) for not using at least two of the following: Abilify Maintena, Abilify Asimtufii, Invega Sustenna, Invega Trinza, or Risperdion Microsphere ER.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

PART B VERSUS PART D

Products Affected

- *acetylcysteine inhalation solution 10 %, 20 %*
- *acyclovir sodium intravenous solution 50 mg/ml*
- *albuterol sulfate inhalation nebulization solution (2.5 mg/3ml) 0.083%, (5 mg/ml) 0.5%, 0.63 mg/3ml, 1.25 mg/3ml, 2.5 mg/0.5ml*
- *amphotericin b intravenous solution reconstituted 50 mg*
- *amphotericin b liposome intravenous suspension reconstituted 50 mg*
- *aprepitant oral capsule 125 mg, 40 mg, 80 & 125 mg, 80 mg*
- **ASTAGRAF XL ORAL CAPSULE EXTENDED RELEASE 24 HOUR 0.5 MG, 1 MG, 5 MG**
- *azathioprine oral tablet 50 mg*
- *budesonide inhalation suspension 0.25 mg/2ml, 0.5 mg/2ml, 1 mg/2ml*
- *clinisol sf intravenous solution 15 %*
- *cromolyn sodium inhalation nebulization solution 20 mg/2ml*
- *cyclophosphamide oral capsule 25 mg, 50 mg*
- *cyclophosphamide oral tablet 25 mg, 50 mg*
- *cyclosporine modified oral capsule 100 mg, 25 mg, 50 mg*
- *cyclosporine modified oral solution 100 mg/ml*
- *cyclosporine oral capsule 100 mg, 25 mg*
- *dronabinol oral capsule 10 mg, 2.5 mg, 5 mg*
- **EMEND ORAL SUSPENSION RECONSTITUTED 125 MG/5ML**
- **ENGERIX-B INJECTION SUSPENSION 20 MCG/ML**
- **ENGERIX-B INJECTION SUSPENSION PREFILLED SYRINGE 10 MCG/0.5ML, 20 MCG/ML**
- **ENVARUSUS XR ORAL TABLET EXTENDED RELEASE 24 HOUR 0.75 MG, 1 MG, 4 MG**
- *everolimus oral tablet 0.25 mg, 0.5 mg, 0.75 mg, 1 mg*
- *formoterol fumarate inhalation nebulization solution 20 mcg/2ml*
- **GAMMAGARD INJECTION SOLUTION 1 GM/10ML, 10 GM/100ML, 2.5 GM/25ML, 20 GM/200ML, 30 GM/300ML, 5 GM/50ML**
- **GAMMAGARD S/D LESS IGA INTRAVENOUS SOLUTION RECONSTITUTED 10 GM, 5 GM**
- **GAMMAKED INJECTION SOLUTION 1 GM/10ML**
- **GAMMAPLEX INTRAVENOUS SOLUTION 10 GM/100ML, 10 GM/200ML, 20 GM/200ML, 5 GM/50ML**
- **GAMUNEX-C INJECTION SOLUTION 1 GM/10ML**
- *gengraf oral capsule 100 mg, 25 mg*
- *gengraf oral solution 100 mg/ml*
- *granisetron hcl oral tablet 1 mg*
- **HEPLISAV-B INTRAMUSCULAR SOLUTION PREFILLED SYRINGE 20 MCG/0.5ML**
- **IMOVAX RABIES INTRAMUSCULAR SUSPENSION RECONSTITUTED 2.5 UNIT/ML**
- **INTRALIPID INTRAVENOUS EMULSION 20 %, 30 %**
- *ipratropium bromide inhalation solution 0.02 %*
- *ipratropium-albuterol inhalation solution 0.5-2.5 (3) mg/3ml*
- *levalbuterol hcl inhalation nebulization solution 0.31 mg/3ml, 0.63 mg/3ml, 1.25 mg/3ml*
- *mycophenolate mofetil oral capsule 250 mg*

- *mycophenolate mofetil oral suspension reconstituted 200 mg/ml*
- *mycophenolate mofetil oral tablet 500 mg*
- *mycophenolate sodium oral tablet delayed release 180 mg, 360 mg*
- *mycophenolic acid oral tablet delayed release 180 mg, 360 mg*
- NULOJIX INTRAVENOUS SOLUTION RECONSTITUTED 250 MG
- NUTRILIPID INTRAVENOUS EMULSION 20 %
- *ondansetron hcl oral solution 4 mg/5ml*
- *ondansetron hcl oral tablet 24 mg, 4 mg, 8 mg*
- *ondansetron oral tablet dispersible 4 mg, 8 mg*
- *pentamidine isethionate inhalation solution reconstituted 300 mg*
- *plenamine intravenous solution 15 %*
- PREHEVBRIO INTRAMUSCULAR SUSPENSION 10 MCG/ML
- PRIVIGEN INTRAVENOUS SOLUTION 10 GM/100ML, 20 GM/200ML, 40 GM/400ML, 5 GM/50ML
- PROGRAF INTRAVENOUS SOLUTION 5 MG/ML
- PROGRAF ORAL PACKET 0.2 MG, 1 MG
- PULMOZYME INHALATION SOLUTION 2.5 MG/2.5ML
- RABAVERT INTRAMUSCULAR SUSPENSION RECONSTITUTED
- RECOMBIVAX HB INJECTION SUSPENSION 10 MCG/ML, 40 MCG/ML, 5 MCG/0.5ML
- RECOMBIVAX HB INJECTION SUSPENSION PREFILLED SYRINGE 10 MCG/ML, 5 MCG/0.5ML
- SANDIMMUNE ORAL SOLUTION 100 MG/ML
- *sirolimus oral solution 1 mg/ml*
- *sirolimus oral tablet 0.5 mg, 1 mg, 2 mg*
- *tacrolimus oral capsule 0.5 mg, 1 mg, 5 mg*
- TENIVAC INTRAMUSCULAR INJECTABLE 5-2 LFU, 5-2 LFU (INJECTION)
- TENIVAC INTRAMUSCULAR SUSPENSION 5-2 LF/0.5ML
- *tobramycin inhalation nebulization solution 300 mg/4ml, 300 mg/5ml*

Details

This drug may be covered under Medicare Part B or D depending upon the circumstances. Information may need to be submitted describing the use and setting of the drug to make the determination.

Index

A

abiraterone acetate 127, 129
 ABIRTEGA 127, 129
 acetylcysteine inhalation solution 10 %, 20
 % 236
 acitretin 1
 ACTEMRA ACTPEN 4
 ACTEMRA SUBCUTANEOUS 4
 ACTHAR 5
 ACTHAR GEL SUBCUTANEOUS PEN-
 INJECTOR..... 5
 acyclovir sodium intravenous solution 50
 mg/ml 236
 ADEMPAS 6
 AIMOVIG..... 26, 27
 AKEEGA 127, 129
 albuterol sulfate inhalation nebulization
 solution (2.5 mg/3ml) 0.083%, (5 mg/ml)
 0.5%, 0.63 mg/3ml, 1.25 mg/3ml, 2.5
 mg/0.5ml 236
 ALECENSA..... 127, 129
 ALUNBRIG..... 127, 129
 ALYFTREK..... 8
 ambrisentan 9
 amphotericin b intravenous solution
 reconstituted 50 mg..... 236
 amphotericin b liposome intravenous
 suspension reconstituted 50 mg 236
 apomorphine hcl subcutaneous 10
 aprepitant oral capsule 125 mg, 40 mg, 80 &
 125 mg, 80 mg 236
 AQNEURSA..... 11
 ARALAST NP INTRAVENOUS
 SOLUTION RECONSTITUTED 1000
 MG, 500 MG..... 7
 ARANESP (ALBUMIN FREE)
 INJECTION SOLUTION 100 MCG/ML,
 200 MCG/ML, 25 MCG/ML, 40
 MCG/ML, 60 MCG/ML 56
 ARANESP (ALBUMIN FREE)
 INJECTION SOLUTION PREFILLED
 SYRINGE 56
 ARCALYST 12

ARIKAYCE..... 13
 ARISTADA INITIO 14
 ARISTADA INTRAMUSCULAR
 PREFILLED SYRINGE 1064
 MG/3.9ML, 441 MG/1.6ML, 662
 MG/2.4ML, 882 MG/3.2ML 14
 armodafinil..... 118
 ASTAGRAF XL ORAL CAPSULE
 EXTENDED RELEASE 24 HOUR 0.5
 MG, 1 MG, 5 MG 236
 AUGTYRO 127, 129
 AUSTEDO 211
 AUSTEDO PATIENT TITRATION KIT
 211
 AUSTEDO XR 211
 AUSTEDO XR PATIENT TITRATION
 ORAL TABLET EXTENDED
 RELEASE THERAPY PACK 12 & 18 &
 24 & 30 MG 211
 AUVELITY 15
 AVMAPKI FAKZYNJA CO-PACK..... 127,
 129
 AYVAKIT 127, 129
 azathioprine oral tablet 50 mg..... 236
B
 bac (butalbital-acetamin-caff)..... 81
 BAFIERTAM 113, 114
 BALVERSA 127, 129
 BENLYSTA SUBCUTANEOUS..... 17
 benztropine mesylate oral 78, 79
 BESREMI 18
 BETASERON SUBCUTANEOUS KIT 113,
 114
 bexarotene 127, 129, 195
 BONSITY 191
 BORUZU 19
 bosentan oral tablet 20
 BOSULIF 127, 129
 BRAFTOVI ORAL CAPSULE 75 MG 127,
 129
 BRUKINSA 127, 129
 budesonide inhalation suspension 0.25
 mg/2ml, 0.5 mg/2ml, 1 mg/2ml 236

butalbital-acetaminophen oral tablet 50-325
 mg 81
 butalbital-apap-caff-cod oral capsule 50-
 325-40-30 mg..... 81
 butalbital-apap-caffeine oral capsule 50-
 325-40 mg 81
 butalbital-apap-caffeine oral solution 81
 butalbital-apap-caffeine oral tablet 50-325-
 40 mg 81
 butalbital-asa-caff-codeine..... 81
 butalbital-aspirin-caffeine oral capsule..... 81
C
 CABOMETYX 127, 129
 CALQUENCE ORAL TABLET ... 127, 129
 CAMCEVI 71
 CAMZYOS 21, 22
 CAPLYTA 130
 CAPRELSA 127, 129
 carglumic acid oral tablet soluble 23
 carisoprodol oral 82
 caspofungin acetate 24
 CAYSTON..... 16
 CERDELGA 25
 chlorzoxazone oral tablet 500 mg 82
 CHOLBAM..... 28
 CIBINQO 29
 CIMZIA (2 SYRINGE) 30, 31
 CIMZIA SUBCUTANEOUS KIT 2 X 200
 MG 30, 31
 CIMZIA-STARTER 30, 31
 CINRYZE 76
 clinisol sf intravenous solution 15 %..... 236
 COBENFY 130
 COBENFY STARTER PACK 130
 COMETRIQ (100 MG DAILY DOSE)
 ORAL KIT 80 & 20 MG 127, 129
 COMETRIQ (140 MG DAILY DOSE)
 ORAL KIT 3 X 20 MG & 80 MG 127,
 129
 COMETRIQ (60 MG DAILY DOSE)... 127,
 129
 COPIKTRA..... 127, 129
 CORLANOR ORAL SOLUTION..... 32
 CORTROPHIN 33
 CORTROPHIN GEL 33
 COSENTYX 34, 35

COSENTYX (300 MG DOSE)..... 34, 35
 COSENTYX SENSOREADY (300 MG) 34,
 35
 COSENTYX SENSOREADY PEN ... 34, 35
 COSENTYX UNOREADY 34, 35
 COTELLIC 127, 129
 cromolyn sodium inhalation nebulization
 solution 20 mg/2ml 236
 CUVRIOR..... 199
 cyclobenzaprine hcl oral tablet 10 mg, 5 mg
 82
 cyclophosphamide oral capsule 25 mg, 50
 mg 236
 cyclophosphamide oral tablet 25 mg, 50 mg
 236
 cyclosporine modified oral capsule 100 mg,
 25 mg, 50 mg 236
 cyclosporine modified oral solution 100
 mg/ml 236
 cyclosporine oral capsule 100 mg, 25 mg236
 cyproheptadine hcl oral 78, 79
 CYSTAGON..... 36
 CYSTARAN 37
D
 dalfampridine er 38
 DANZITEN 127, 129
 dasatinib 127, 129
 DAURISMO 127, 129
 deferasirox..... 39
 deferasirox granules 39
 deferiprone 40
 DIACOMIT..... 41
 dichlorphenamide..... 42
 DIFICID 43
 dihydroergotamine mesylate nasal..... 44
 dimethyl fumarate oral capsule delayed
 release 120 mg, 240 mg 113, 114
 dimethyl fumarate starter pack oral capsule
 delayed release therapy pack 113, 114
 diphenoxylate-atropine oral liquid..... 78, 79
 diphenoxylate-atropine oral tablet 2.5-0.025
 mg 78, 79
 dipyrindamole oral 78, 79
 DOPTELET 45
 DOPTELET SPRINKLE 45
 doxepin hcl external 46

dronabinol oral capsule 10 mg, 2.5 mg, 5 mg
..... 236

DUPIXENT SUBCUTANEOUS
SOLUTION AUTO-INJECTOR 47, 48

DUPIXENT SUBCUTANEOUS
SOLUTION PREFILLED SYRINGE . 47,
48

E

EGRIFTA SV..... 49

EGRIFTA WR 49

ELIGARD 71

eltrombopag olamine oral packet 12.5 mg,
25 mg 151

eltrombopag olamine oral tablet 12.5 mg, 25
mg, 50 mg, 75 mg 151

EMEND ORAL SUSPENSION
RECONSTITUTED 125 MG/5ML 236

EMGALITY 26, 27

EMGALITY (300 MG DOSE) 26, 27

EMSAM..... 50

ENBREL MINI..... 51

ENBREL SUBCUTANEOUS SOLUTION
25 MG/0.5ML 51

ENBREL SUBCUTANEOUS SOLUTION
PREFILLED SYRINGE 51

ENBREL SURECLICK SUBCUTANEOUS
SOLUTION AUTO-INJECTOR 51

ENGERIX-B INJECTION SUSPENSION
20 MCG/ML 236

ENGERIX-B INJECTION SUSPENSION
PREFILLED SYRINGE 10 MCG/0.5ML,
20 MCG/ML 236

ENTYVIO PEN 53

ENVARUSUS XR ORAL TABLET
EXTENDED RELEASE 24 HOUR 0.75
MG, 1 MG, 4 MG 236

EPIDIOLEX..... 54

EPOGEN INJECTION SOLUTION 10000
UNIT/ML, 2000 UNIT/ML, 20000
UNIT/ML, 3000 UNIT/ML, 4000
UNIT/ML 56

ergotamine-caffeine 78, 79

ERIVEDGE..... 127, 129

ERLEADA..... 127, 129

erlotinib hcl 127, 129

ERZOFRI INTRAMUSCULAR
SUSPENSION PREFILLED SYRINGE
117 MG/0.75ML, 156 MG/ML, 234
MG/1.5ML, 351 MG/2.25ML, 39
MG/0.25ML, 78 MG/0.5ML 57

estradiol oral..... 80

estradiol transdermal patch twice weekly . 80

estradiol transdermal patch weekly..... 80

eszopiclone..... 83

EUCRISA 58

EULEXIN 127, 129

everolimus oral tablet 0.25 mg, 0.5 mg, 0.75
mg, 1 mg 236

everolimus oral tablet 10 mg, 2.5 mg, 5 mg,
7.5 mg 127, 129

everolimus oral tablet soluble 127, 129

EVRYSDI 59

F

FABHALTA 60

FANAPT 130

FANAPT TITRATION PACK A 130

FANAPT TITRATION PACK B ORAL
TABLET 130

FANAPT TITRATION PACK C ORAL
TABLET 130

FASENRA 61

FASENRA PEN..... 61

fentanyl citrate buccal lozenge on a handle
..... 62

fidaxomicin 43

FILSPARI 63

figolimod hcl 113, 114

FINTEPLA..... 64

FIRDAPSE..... 65

FIRMAGON (240 MG DOSE)..... 71

FIRMAGON SUBCUTANEOUS
SOLUTION RECONSTITUTED 80 MG
..... 71

flucytosine oral..... 66

fluorouracil external cream 0.5 %..... 67

formoterol fumarate inhalation nebulization
solution 20 mcg/2ml 236

FOTIVDA 127, 129

FRUZAQLA 127, 129

FULPHILA 217

FYLNETRA..... 217

G

GALAFOLD	68
GAMMAGARD INJECTION SOLUTION	
1 GM/10ML, 10 GM/100ML, 2.5	
GM/25ML, 20 GM/200ML, 30	
GM/300ML, 5 GM/50ML	236
GAMMAGARD S/D LESS IGA	
INTRAVENOUS SOLUTION	
RECONSTITUTED 10 GM, 5 GM....	236
GAMMAKED INJECTION SOLUTION 1	
GM/10ML	236
GAMMAPLEX INTRAVENOUS	
SOLUTION 10 GM/100ML, 10	
GM/200ML, 20 GM/200ML, 5	
GM/50ML	236
GAMUNEX-C INJECTION SOLUTION 1	
GM/10ML	236
GATTEX.....	69
GAVRETO	127, 129
gefitinib	127, 129
gengraf oral capsule 100 mg, 25 mg	236
gengraf oral solution 100 mg/ml	236
GENOTROPIN MINIQUICK	
SUBCUTANEOUS PREFILLED	
SYRINGE	73, 74
GENOTROPIN SUBCUTANEOUS	
CARTRIDGE.....	73, 74
GILOTRIF	127, 129
GLASSIA.....	7
glatiramer acetate subcutaneous solution	
prefilled syringe 20 mg/ml, 40 mg/ml	113, 114
glatopa subcutaneous solution prefilled	
syringe 20 mg/ml, 40 mg/ml	113, 114
glyburide micronized oral tablet 1.5 mg, 3	
mg, 6 mg	78, 79
glyburide oral tablet 1.25 mg, 2.5 mg, 5 mg	
.....	78, 79
glyburide-metformin oral tablet 1.25-250	
mg, 2.5-500 mg	78, 79
glyburide-metformin oral tablet 5-500 mg	
.....	78, 79
GOCOVRI	72
GOMEKLI	127, 129
granisetron hcl oral tablet 1 mg	236
guanfacine hcl er	78, 79

guanfacine hcl oral	78, 79
---------------------------	--------

H

HADLIMA.....	75
HADLIMA PUSHTOUCH.....	75
HAEGARDA	76
HEPLISAV-B INTRAMUSCULAR	
SOLUTION PREFILLED SYRINGE 20	
MCG/0.5ML	236
HERNEXEOS.....	127, 129
HETLIOZ LQ	77
HUMATROPE INJECTION CARTRIDGE	
.....	73, 74
HUMIRA (1 PEN)	84, 85
HUMIRA (2 PEN) SUBCUTANEOUS	
AUTO-INJECTOR KIT.....	84, 85
HUMIRA (2 SYRINGE)	
SUBCUTANEOUS PREFILLED	
SYRINGE KIT 10 MG/0.1ML, 20	
MG/0.2ML, 40 MG/0.4ML, 40	
MG/0.8ML	84, 85
HUMIRA-CD/UC/HS STARTER	
SUBCUTANEOUS AUTO-INJECTOR	
KIT 80 MG/0.8ML	84, 85
HUMIRA-PED>=40KG UC STARTER	
SUBCUTANEOUS AUTO-INJECTOR	
KIT	84, 85
HUMIRA-PSORIASIS/UEVIT STARTER	
SUBCUTANEOUS AUTO-INJECTOR	
KIT	84, 85
hydroxyzine hcl oral syrup.....	78, 79
hydroxyzine hcl oral tablet 25 mg, 50 mg	78, 79
hydroxyzine pamoate oral	78, 79
HYFTOR.....	86

I

IBRANCE	127, 129
IBTROZI.....	127, 129
icatibant acetate subcutaneous solution	
prefilled syringe	87
ICLUSIG.....	127, 129
IDHIFA	127, 129
ILARIS SUBCUTANEOUS SOLUTION	88
ILUMYA.....	89
imatinib mesylate oral.....	127, 129
IMBRUVICA ORAL CAPSULE	90

IMBRUVICA ORAL SUSPENSION.... 127, 129
IMBRUVICA ORAL TABLET 140 MG, 280 MG, 420 MG..... 90
IMKELDI..... 127, 129
IMOVAX RABIES INTRAMUSCULAR SUSPENSION RECONSTITUTED 2.5 UNIT/ML 236
IMPAVIDO..... 91
INCRELEX..... 92
indomethacin oral capsule 25 mg, 50 mg 78, 79
INGREZZA ORAL CAPSULE..... 211
INGREZZA ORAL CAPSULE SPRINKLE 211
INGREZZA ORAL CAPSULE THERAPY PACK..... 211
INLYTA..... 127, 129
INQOVI 127, 129
INREBIC..... 127, 129
INTRALIPID INTRAVENOUS EMULSION 20 %, 30 % 236
ipratropium bromide inhalation solution 0.02 % 236
ipratropium-albuterol inhalation solution 0.5-2.5 (3) mg/3ml 236
ITOVEBI..... 127, 129
ivabradine hcl..... 32
IWILFIN 127, 129
J
JAKAFI..... 93
JAYPIRCA 127, 129
JYLAMVO 94
K
KALYDECO..... 95
KERENDIA 96
KESIMPTA..... 113, 114
ketorolac tromethamine oral 78, 79
KEVZARA 97
KINERET SUBCUTANEOUS SOLUTION PREFILLED SYRINGE 98
KISQALI (200 MG DOSE) 127, 129
KISQALI (400 MG DOSE) 127, 129
KISQALI (600 MG DOSE) 127, 129
KISQALI FEMARA (200 MG DOSE) . 127, 129

KISQALI FEMARA (400 MG DOSE) . 127, 129
KISQALI FEMARA (600 MG DOSE) . 127, 129
KOSELUGO ORAL CAPSULE 127, 129
KRAZATI..... 127, 129
L
lapatinib ditosylate 127, 129
LAZCLUZE..... 127, 129
lenalidomide..... 127, 129
LENVIMA (10 MG DAILY DOSE) 127, 129
LENVIMA (12 MG DAILY DOSE) 127, 129
LENVIMA (14 MG DAILY DOSE) 127, 129
LENVIMA (18 MG DAILY DOSE) 127, 129
LENVIMA (20 MG DAILY DOSE) 127, 129
LENVIMA (24 MG DAILY DOSE) 127, 129
LENVIMA (4 MG DAILY DOSE) 127, 129
LENVIMA (8 MG DAILY DOSE) 127, 129
LEQSELVI 99
LEUKERAN 127, 129
LEUKINE INJECTION SOLUTION RECONSTITUTED..... 217
leuprolide acetate (3 month) 71
levalbuterol hcl inhalation nebulization solution 0.31 mg/3ml, 0.63 mg/3ml, 1.25 mg/3ml 236
l-glutamine oral packet 52
LIBERVANT..... 100
lidocaine external patch 5 % 197
LITFULO 101
LIVMARLI..... 102
LIVTENCITY..... 103
LODOCO 104
lofexidine hcl 105
LONSURF 127, 129
LORBRENA 127, 129
LUMAKRAS 127, 129
LUPKYNIS..... 106
LUPRON DEPOT (1-MONTH)..... 71
LUPRON DEPOT (3-MONTH)..... 71

LUPRON DEPOT (4-MONTH) 71
 LUPRON DEPOT (6-MONTH) 71
 LUTRATE DEPOT 71
 LYBALVI 107
 LYNPARZA ORAL TABLET 127, 129
 LYTGObI (12 MG DAILY DOSE) 127, 129
 LYTGObI (16 MG DAILY DOSE) 127, 129
 LYTGObI (20 MG DAILY DOSE) 127, 129
M
 MAVENCLAD (10 TABS) 113, 114
 MAVENCLAD (4 TABS) 113, 114
 MAVENCLAD (5 TABS) 113, 114
 MAVENCLAD (6 TABS) 113, 114
 MAVENCLAD (7 TABS) 113, 114
 MAVENCLAD (8 TABS) 113, 114
 MAVENCLAD (9 TABS) 113, 114
 MAVYRET 108
 MAYZENT 113, 114
 MAYZENT STARTER PACK 113, 114
 megestrol acetate oral suspension 78, 79
 megestrol acetate oral tablet 80
 MEKINIST 127, 129
 MEKTOVI 127, 129
 MENEST 80
 mercaptopurine oral suspension 127, 129
 metaxalone oral tablet 800 mg 82
 methocarbamol oral tablet 500 mg, 750 mg
 82
 methyltestosterone oral 109
 metyrosine 110
 mifepristone oral tablet 300 mg 111
 miglustat 112
 modafinil oral 118
 MODEYSO 127, 129
 MOUNJARO SUBCUTANEOUS
 SOLUTION AUTO-INJECTOR 70
 mycophenolate mofetil oral capsule 250 mg
 236
 mycophenolate mofetil oral suspension
 reconstituted 200 mg/ml 237
 mycophenolate mofetil oral tablet 500 mg
 237
 mycophenolate sodium oral tablet delayed
 release 180 mg, 360 mg 237
 mycophenolic acid oral tablet delayed
 release 180 mg, 360 mg 237

MYFEMBREE 115
N
 NAYZILAM 116
 NERLYNX 128, 129
 NEULASTA ONPRO 217
 NEULASTA SUBCUTANEOUS
 SOLUTION PREFILLED SYRINGE 217
 NEXLETOL 2, 3
 NEXLIZET 2, 3
 NGENLA 73, 74
 nifedipine oral 78, 79
 nilotinib d-tartrate 128, 129
 nilotinib hcl 128, 129
 nilutamide 128, 129
 NINLARO 128, 129
 nitisinone 117
 NIVESTYM 217
 NORDITROPIN FLEXPRO
 SUBCUTANEOUS SOLUTION PEN-
 INJECTOR 73, 74
 NUBEQA 128, 129
 NUCALA 119
 NUEDEXTA 120
 NULOJIX INTRAVENOUS SOLUTION
 RECONSTITUTED 250 MG 237
 NUPLAZID ORAL CAPSULE 121
 NUPLAZID ORAL TABLET 10 MG 121
 NURTEC 26, 27
 NUTRILIPID INTRAVENOUS
 EMULSION 20 % 237
 NUTROPIN AQ NUSPIN 10
 SUBCUTANEOUS SOLUTION PEN-
 INJECTOR 73, 74
 NUTROPIN AQ NUSPIN 20
 SUBCUTANEOUS SOLUTION PEN-
 INJECTOR 73, 74
 NUTROPIN AQ NUSPIN 5
 SUBCUTANEOUS SOLUTION PEN-
 INJECTOR 73, 74
 NYVEPRIA 217
O
 OCREVUS ZUNOVO 122
 octreotide acetate injection solution 100
 mcg/ml, 1000 mcg/ml, 200 mcg/ml, 50
 mcg/ml, 500 mcg/ml 123
 octreotide acetate intramuscular 123

ODOMZO 128, 129
 OFEV 124
 OGSIVEO 128, 129
 OJEMDA 128, 129
 OJJAARA 128, 129
 OMNITROPE SUBCUTANEOUS
 SOLUTION CARTRIDGE..... 73, 74
 OMNITROPE SUBCUTANEOUS
 SOLUTION RECONSTITUTED... 73, 74
 ondansetron hcl oral solution 4 mg/5ml.. 237
 ondansetron hcl oral tablet 24 mg, 4 mg, 8
 mg 237
 ondansetron oral tablet dispersible 4 mg, 8
 mg 237
 ONUREG..... 128, 129
 OPDIVO QVANTIG 125
 OPIPZA ORAL FILM 10 MG, 2 MG, 5 MG
 130
 OPSUMIT 126
 ORENCIA CLICKJECT..... 131
 ORENCIA INTRAVENOUS 131
 ORENCIA SUBCUTANEOUS SOLUTION
 PREFILLED SYRINGE 131
 ORFADIN ORAL SUSPENSION..... 117
 ORGOVYX..... 128, 129
 ORIAHNN 132
 ORILISSA..... 133
 ORKAMBI..... 134
 ORLADEYO..... 76
 orphenadrine citrate er 82
 ORSERDU..... 128, 129
 OSENVLT..... 222
 OTEZLA 135
 OTEZLA XR 135
 OTEZLA/OTEZLA XR INITIATION PK
 135
 OXERVATE 136
 oxycodone hcl er oral tablet er 12 hour
 abuse-deterrent 10 mg, 20 mg, 40 mg, 80
 mg 137, 138
 OXYCONTIN ORAL TABLET ER 12
 HOUR ABUSE-DETERRENT 10 MG,
 15 MG, 20 MG, 30 MG, 40 MG, 60 MG,
 80 MG 137, 138

OZEMPIC (0.25 OR 0.5 MG/DOSE)
 SUBCUTANEOUS SOLUTION PEN-
 INJECTOR 2 MG/3ML 70
 OZEMPIC (1 MG/DOSE)
 SUBCUTANEOUS SOLUTION PEN-
 INJECTOR 4 MG/3ML 70
 OZEMPIC (2 MG/DOSE) 70
P
 PANRETIN..... 195
 pazopanib hcl 128, 129
 PEGASYS SUBCUTANEOUS SOLUTION
 180 MCG/ML 141
 PEGASYS SUBCUTANEOUS SOLUTION
 PREFILLED SYRINGE 141
 PEMAZYRE..... 128, 129
 penicillamine oral tablet..... 142
 pentamidine isethionate inhalation solution
 reconstituted 300 mg..... 237
 pentamidine isethionate injection 143
 perphenazine-amitriptyline 80
 PERSERIS 144
 phenobarbital oral elixir 20 mg/5ml 80
 phenobarbital oral tablet 80
 phenoxybenzamine hcl oral 145
 PIASKY 146
 PIQRAY (200 MG DAILY DOSE) 128, 129
 PIQRAY (250 MG DAILY DOSE) 128, 129
 PIQRAY (300 MG DAILY DOSE) 128, 129
 pirfenidone 147
 plenamine intravenous solution 15 %..... 237
 POMALYST 128, 129
 PONVORY 113, 114
 PONVORY STARTER PACK..... 113, 114
 posaconazole oral..... 148
 PRALUENT SUBCUTANEOUS
 SOLUTION AUTO-INJECTOR 139, 140
 PREHEVBRIO INTRAMUSCULAR
 SUSPENSION 10 MCG/ML 237
 PREMARIN ORAL 80
 PRETOMANID 149
 PREVYMIS ORAL 150
 PRIVIGEN INTRAVENOUS SOLUTION
 10 GM/100ML, 20 GM/200ML, 40
 GM/400ML, 5 GM/50ML 237
 PROCRT..... 56

PROGRAF INTRAVENOUS SOLUTION 5
 MG/ML 237
 PROGRAF ORAL PACKET 0.2 MG, 1 MG
 237
 PROLASTIN-C INTRAVENOUS
 SOLUTION..... 7
 promethazine hcl oral solution 6.25 mg/5ml
 78, 79
 promethazine hcl oral syrup..... 78, 79
 promethazine hcl oral tablet..... 78, 79
 promethazine hcl rectal suppository 12.5
 mg, 25 mg 78, 79
 promethazine-phenylephrine..... 78, 79
 promethegan rectal suppository 50 mg 78, 79
 PULMOZYME INHALATION
 SOLUTION 2.5 MG/2.5ML 237
 PYRUKYND 152
 PYRUKYND TAPER PACK..... 152
Q
 QINLOCK..... 128, 129
 QULIPTA 26, 27
R
 RABAVERT INTRAMUSCULAR
 SUSPENSION RECONSTITUTED... 237
 RADICAVA ORS..... 153
 RADICAVA ORS STARTER KIT 153
 RAVICTI 154
 REBIF REBIDOSE SUBCUTANEOUS
 SOLUTION AUTO-INJECTOR 113, 114
 REBIF REBIDOSE TITRATION PACK
 SUBCUTANEOUS SOLUTION AUTO-
 INJECTOR..... 113, 114
 REBIF SUBCUTANEOUS SOLUTION
 PREFILLED SYRINGE 113, 114
 REBIF TITRATION PACK
 SUBCUTANEOUS SOLUTION
 PREFILLED SYRINGE 113, 114
 RECOMBIVAX HB INJECTION
 SUSPENSION 10 MCG/ML, 40
 MCG/ML, 5 MCG/0.5ML 237
 RECOMBIVAX HB INJECTION
 SUSPENSION PREFILLED SYRINGE
 10 MCG/ML, 5 MCG/0.5ML 237
 RECORLEV 155
 RELEUKO SUBCUTANEOUS..... 217
 RELISTOR ORAL..... 156

RELISTOR SUBCUTANEOUS
 SOLUTION..... 156
 RELISTOR SUBCUTANEOUS
 SOLUTION PREFILLED SYRINGE 156
 REPATHA 139, 140
 REPATHA SURECLICK..... 139, 140
 RETACRIT INJECTION SOLUTION
 10000 UNIT/ML, 10000
 UNIT/ML(1ML), 2000 UNIT/ML, 20000
 UNIT/ML, 3000 UNIT/ML, 4000
 UNIT/ML, 40000 UNIT/ML 56
 RETEVMO 128, 129
 REVLIMID 128, 129
 REVUFORJ 128, 129
 REXULTI 157
 REZLIDHIA 128, 129
 REZUROCK..... 158
 RINVOQ..... 159, 160
 RINVOQ LQ..... 159, 160
 ROMVIMZA 128, 129
 ROZLYTREK..... 128, 129
 RUBRACA 128, 129
 rufinamide oral suspension 161
 rufinamide oral tablet..... 161
 RYBELSUS 70
 RYBELSUS (FORMULATION R2)..... 70
 RYDAPT..... 128, 129
 RYKINDO 162
 RYLAZE..... 163
S
 SANDIMMUNE ORAL SOLUTION 100
 MG/ML 237
 sapropterin dihydrochloride oral packet . 164
 sapropterin dihydrochloride oral tablet... 164
 SCEMBLIX 128, 129
 SECUADO..... 165
 SEROSTIM SUBCUTANEOUS
 SOLUTION RECONSTITUTED 4 MG,
 5 MG, 6 MG..... 166
 SIGNIFOR 167
 sildenafil citrate oral suspension
 reconstituted..... 168
 sildenafil citrate oral tablet 20 mg 168
 SILIQ 169
 SIMPONI SUBCUTANEOUS SOLUTION
 AUTO-INJECTOR 170

SIMPONI SUBCUTANEOUS SOLUTION	
PREFILLED SYRINGE	170
sirolimus oral solution 1 mg/ml	237
sirolimus oral tablet 0.5 mg, 1 mg, 2 mg	237
SIRTURO	171
SKYRIZI	172
SKYRIZI PEN	172
SKYTROFA	73, 74
sodium oxybate	173
sodium phenylbutyrate oral powder 3 gm/tsp	174
sodium phenylbutyrate oral tablet	174
SOFOSBUVIR-VELPATASVIR	175
SOLTAMOX	128, 129
SOMAVERT	176
sorafenib tosylate	128, 129
SOTYKTU	177
STELARA INTRAVENOUS	178
STELARA SUBCUTANEOUS	
SOLUTION 45 MG/0.5ML	178
STELARA SUBCUTANEOUS	
SOLUTION PREFILLED SYRINGE	178
STIMUFEND	217
STIVARGA	128, 129
SUCRAID	179
sunitinib malate	128, 129
SYMDEKO	180
SYMLINPEN 120 SUBCUTANEOUS	
SOLUTION PEN-INJECTOR	181
SYMLINPEN 60 SUBCUTANEOUS	
SOLUTION PEN-INJECTOR	181
SYNAREL	182
T	
TABLOID	128, 129
TABRECTA	128, 129
tacrolimus oral capsule 0.5 mg, 1 mg, 5 mg	237
tadalafil (pah)	183
tadalafil oral tablet 5 mg	184
TADLIQ	183
TAFINLAR	128, 129
TAGRISSO	128, 129
TALTZ	185
TALZENNA	128, 129
TARPEYO	186
TASCENSO ODT	113, 114
tasimelteon	77
TAVNEOS	187
TAZVERIK	128, 129
TECENTRIQ HYBREZA	188
TEFLARO	189
temazepam	83
TENIVAC INTRAMUSCULAR	
INJECTABLE 5-2 LFU, 5-2 LFU	
(INJECTION)	237
TENIVAC INTRAMUSCULAR	
SUSPENSION 5-2 LF/0.5ML	237
TEPEZZA	190
TEPMETKO	128, 129
teriflunomide	113, 114
TERIPARATIDE SUBCUTANEOUS	
SOLUTION PEN-INJECTOR 560	
MCG/2.24ML	191
testosterone transdermal gel 1.62 %, 12.5	
mg/act (1%), 20.25 mg/1.25gm (1.62%),	
20.25 mg/act (1.62%), 25 mg/2.5gm	
(1%), 40.5 mg/2.5gm (1.62%), 50	
mg/5gm (1%)	196
testosterone transdermal solution	196
tetrabenazine	211
THALOMID	128, 129
TIBSOVO	128, 129
tigecycline	193
tiopronin oral	192
tobramycin inhalation nebulization solution	
300 mg/4ml, 300 mg/5ml	237
tolvaptan	194
topiramate oral solution	55
toremifene citrate	128, 129
TRELSTAR MIXJECT	71
TREMFYA CROHNS INDUCTION	198
TREMFYA ONE-PRESS	
SUBCUTANEOUS SOLUTION PEN-	
INJECTOR	198
TREMFYA PEN	198
TREMFYA SUBCUTANEOUS	
SOLUTION PREFILLED SYRINGE	198
TREMFYA-CD/UC INDUCTION	198
tretinoin oral	128, 129
trientine hcl oral capsule 250 mg	199
trihexyphenidyl hcl	78, 79
TRIKAFTA	200

TRULICITY SUBCUTANEOUS
 SOLUTION AUTO-INJECTOR 70
 TRUQAP..... 128, 129
 TUKYSA 128, 129
 TURALIO ORAL CAPSULE 125 MG. 128,
 129
 TYMLOS 201
 TYVASO DPI MAINTENANCE KIT
 INHALATION POWDER 16 MCG, 32
 MCG, 48 MCG, 64 MCG 202
 TYVASO DPI TITRATION KIT
 INHALATION POWDER 16 & 32 & 48
 MCG 202
U
 UBRELVY..... 26, 27
 UDENYCA 217
 UDENYCA ONBODY 217
 UPTRAVI ORAL 203
 UPTRAVI TITRATION..... 203
 UZEDY 204
V
 VALCHLOR..... 205
 VALTOCO 10 MG DOSE..... 116
 VALTOCO 15 MG DOSE NASAL
 LIQUID THERAPY PACK 2 X 7.5
 MG/0.1ML 116
 VALTOCO 20 MG DOSE NASAL
 LIQUID THERAPY PACK 2 X 10
 MG/0.1ML 116
 VALTOCO 5 MG DOSE..... 116
 VANFLYTA 128, 129
 VEMLIDY 206
 VENCLEXTA..... 128, 129
 VENCLEXTA STARTING PACK 128, 129
 VENTAVIS..... 207
 VEOZAH 208
 VERZENIO..... 128, 129
 VICTOZA SUBCUTANEOUS SOLUTION
 PEN-INJECTOR..... 70
 vigabatrin 209
 VIGAFYDE 209
 VIJOICE 210
 VITRAKVI 128, 129
 VIZIMPRO 128, 129
 VONJO 128, 129
 VORANIGO 128, 129

voriconazole intravenous 212
 VOSEVI..... 213
 VOWST 214
 VRAYLAR ORAL CAPSULE..... 130
W
 WEGOVY SUBCUTANEOUS SOLUTION
 AUTO-INJECTOR 0.25 MG/0.5ML, 0.5
 MG/0.5ML, 1 MG/0.5ML, 1.7
 MG/0.75ML, 2.4 MG/0.75ML ... 215, 216
 WELIREG..... 128, 129
 WYOST 222
X
 XALKORI..... 128, 129
 XATMEP 218
 XDEMVI 219
 XELJANZ 220
 XELJANZ XR 220
 XERMELO 221
 XGEVA..... 222
 XIFAXAN..... 223
 XOLAIR 224, 225
 XOLREMDI 226
 XOSPATA 128, 129
 XPOVIO (100 MG ONCE WEEKLY)
 ORAL TABLET THERAPY PACK 50
 MG 128, 129
 XPOVIO (40 MG ONCE WEEKLY) ORAL
 TABLET THERAPY PACK 10 MG, 40
 MG 128, 129
 XPOVIO (40 MG TWICE WEEKLY)
 ORAL TABLET THERAPY PACK 40
 MG 128, 129
 XPOVIO (60 MG ONCE WEEKLY) ORAL
 TABLET THERAPY PACK 60 MG. 128,
 129
 XPOVIO (60 MG TWICE WEEKLY).. 128,
 129
 XPOVIO (80 MG ONCE WEEKLY) ORAL
 TABLET THERAPY PACK 40 MG. 128,
 129
 XPOVIO (80 MG TWICE WEEKLY).. 128,
 129
 XTANDI 128, 129
 XYWAV 227
Y
 YONSA..... 128, 129

YORVIPATH	228
YUTREPIA.....	229
Z	
zaleplon	83
ZARXIO	217
ZAVZPRET	26, 27
ZEJULA ORAL TABLET.....	128, 129
ZELBORAF	128, 129
ZEMAIRA	7
ZEPBOUND SUBCUTANEOUS SOLUTION AUTO-INJECTOR	230
ZEPOSIA	231
ZEPOSIA 7-DAY STARTER PACK.....	231
ZEPOSIA STARTER KIT ORAL CAPSULE THERAPY PACK 0.23MG &0.46MG 0.92MG(21).....	231

ZIEXTENZO	217
ZILBRYSQ	232
ZOLINZA	128, 129
zolpidem tartrate er	83
zolpidem tartrate oral tablet 10 mg	83
ZTALMY	233
ZTLIDO	197
ZURZUVAE	234
ZYDELIG	128, 129
ZYKADIA ORAL TABLET	128, 129
ZYPREXA RELPREVV INTRAMUSCULAR SUSPENSION RECONSTITUTED 210 MG, 300 MG, 405 MG	235